



“Treatment and interference of Common Complications of Chronic Kidney disorder: A study in Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh”

Dr. Md Abdur Rouf^{1*}, AKM Khairul Islam², Dr. Md Atiqul Islam³

¹ Registrar, Renal & Dialysis Unit, Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh

² RMO, Emergency, Observation and Referral Unit, Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh

³ Assistant Professor, Department of Paediatric Infectious Diseases and Community Paediatrics, Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh

Abstract

Chronic Kidney disorder (CKD) could be a worldwide public ill health with associate increasing incidence and prevalence. Outcomes of CKD embrace not solely complications of small excretory organ perform and disorder however additionally nephrosis inflicting hyperbolic morbidity and mortality. Sadly, CKD is commonly undetected and undertreated as a result of its insidious onset, variable progression, and length of time to explicit nephrosis. This study was conduct June 2016 form July 2017 serve Renal & Dialysis Unit Dhaka shishu (Children) Hospital a damage or decreased kidney polygenic disorder is currently the leading reason behind CKD requiring excretory organ replacement medical aid in several elements of the globe, and its prevalence is increasing disproportionately within the developing countries. This literary criticism outlines the present recommendations from numerous clinical pointers and analysis studies for treatment, hindrance and delaying the progression of each CKD and its common complications like vital sign cardiovascular disease, anemia, excretory organ dystrophy, solution and acid base imbalance. Recommendations for nutrition in CKD and measures adopted for early diabetic nephrosis to stop any progression have additionally been reviewed. This study proof that early detection and management of CKD will forestall or cut back illness progression, decrease complications and improve outcomes proof supports that achieving best aldohexose management, blood pressure, reduction in symptom with a complex intervention slows the progression of CKD. Angiotensin-converting catalyst inhibitors and angiotensin-2 receptor antagonists are best as a result of their distinctive ability to decrease symptom, an element necessary for the progression of CKD.

Keywords: chronic kidney disease, diabetic kidney disease, end stage renal disease

1. Introduction

Chronic uropathy could be a major international public pathological state equally moving the folks of developed countries likewise as developing countries. it's recognized as a risk issue for disorder (CVD), cardiovascular disease, excretory organ dystrophy, anemia and deficiency disease. ^[1] once advanced, it additionally carries a better risk of mortality. ^[2] As excretory organ perform deteriorates, patients develop complications associated with fluid overload, solution and acid-base imbalances, and also the build-up of element waste product. To survive, some patients eventually would like dialysis or excretory organ transplantation. However treatment with qualitative analysis or transplantation is extremely big-ticket and troublesome for peoples of nations like Peoples of Bangladesh. CKD is often unrecognized and is commonly undertreated. Early recognition and intervention area unit essential to slow sickness progression, maintain quality of life and improve outcome. It's necessary to spot the reason behind CKD so specific medical aid will be instituted. Causes of CKD embody nephritis, DM (DM), cardiovascular disease, tract infection, tract obstruction, tubulointerstitial nephrosis and infiltrative diseases. Studies in South Asian countries together with People's Republic of Bangladesh have shown glomerulonephritis nephritis Bright's sickness to be the foremost common reason behind finish stage excretory organ disease (ESRD) followed by DM. ^[3-5] the foremost

frequent reason behind CKD in North America and Europe is DM followed by cardiovascular disease. ⁶ DM is currently turning into more and more rife in developing countries, maybe because of amendment in dietary habits, diminished physical activity and weight gain ^[6]. so, DM is currently the leading reason behind ESRD requiring excretory organ replacement medical aid (RRT) in several components of the globe, and its prevalence is increasing disproportionately within the developing world. ⁶ Patients with fresh diagnosed CKD usually additionally gift with cardiovascular disease. once no bald proof for a primary capillary or tubulointerstitial uropathy method is gift, CKD is commonly attributed to cardiovascular disease ^[6]. The incidence of ESRD in our country isn't famous, however would be a lot of above that in developed countries thanks to high incidence of infection and environmental pollution likewise as increasing prevalence of DM. this text critically reviews this recommendations and therapeutic methods revealed in recent journals and numerous clinical tips for treatment and preventing or delaying the progression of CKD and it's common complications.

2. Method and Materials

This study was conduct June 2016 form July 2017 serve the Renal & Dialysis Unit, Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh a damage or decreased kidney function for 3 months or more. The definition of CKD is the same

regardless of age, sex and race. This definition is widely accepted and has been adopted in various clinical guidelines.^{2, 8-13} Decreased kidney function means a glomerular filtration rate (GFR) of less than 60 mL per minute per 1.73 m² body surface area ^[7]. A GFR rate of more than 60 mL/min/1.73 m² is considered abnormal if it is accompanied by documented evidence of kidney damage. The principal marker of urinary organ injury is that the presence of persistent macromolecule in pee. Alternative proof of urinary organ injury includes abnormalities in pee sediment, blood and pee organic chemistry, or abnormal findings in imaging studies ^[7]. CKD may be outlined by the presence of urinary simple protein in associate degree excretion rate on top of three hundred mg per twenty four hours macro albuminuria or urinary protein excretion 0.5 g/24 hours or more buminuria or 24-hour urinary albumin excretion in the range of 30 to 300 mg indicates the early stage of kidney disease in DM.¹⁵

3. Results

The GFR is that the best live of overall urinary organ perform ^[7]. the traditional level of GFR varies per age, sex, and body size. In healthy young adult of thirty years or younger, the traditional GFR is close to one hundred twenty five mL/min per 1.73 money supply and declines by close to one mL/min per 1.73 money supply each year thenceforth.⁷ Estimation of the GFR not needs a 24-hour excrement assortment for creatinine clearance; it will be accomplished with similar accuracy victimization mathematic formulas. Most commonly used formulas are the Modification of Diet in Renal Disease (MDRD) equation and the Cockcroft-Gault equation (Table I). Many laboratories now report an estimated GFR (eGFR), using one of these equations. The MDRD has the advantage of being more accurate than the Cockcroft-Gault in persons with a GFR less than 90 mL/min per 1.73 m² and takes race into consideration.^{7, 16, 17} At GFR below 60 mL/min, the prevalence of complications of CKD increases.¹⁸⁻²⁰ However, in people aged over 12 years, an eGFR in the range 45–59 mL/min/1.73 m², if stable over time and without any other evidence of kidney damage, is unlikely to be associated with CKD related complications.²

Table 1: Recommended equations for estimation of GFR.

Equation from the Modification of Diet in Renal
Disease study
Estimated GFR (mL/min per 1.73 m ²)
$186 \times (S. Cr \text{ mg/dL})^{-1.154} \times (\text{age in years})^{-0.203}$
Multiply by 0.742 for women
Multiply by 1.21 for African Americans
Cockcroft Gault equation
Estimated creatinine clearance (mL/min)
$(140 - \text{age in years}) \times \text{Body weight (kg)}$
$= 72 \times S. Cr \text{ (mg/dL)}$
Multiply by 0.85 for women

Tags of CKD

The NKF classifies CKD into five stages (Table 2) based on the GFR and/or presence of kidney damage.⁷ This staging system is widely accepted by various clinical guidelines on the management of chronic kidney disease.^{2,8-13} excreted by the kidneys results in the uremic syndrome. This syndrome results in death unless the toxins ar removed by RRT, victimization chemical analysis or urinary organ transplantation ^[6]., twenty two Denial of medical care means that death in days, weeks, or months. ESRD includes patients treated by chemical analysis or transplantation, regardless of the extent of GFR ^[7].

Table 2: Classification of chronic kidney disease

Stage	GFR (mL/min per 1.73 m ²)	Description
1	##	Kidney damage with normal or GFR
2	10–12	Kidney damage with Mild GFR
3	5–10	Moderate GFR
4	1–5	Severe GFR
5	<1	Kidney failure

Management Maintenance of euvolemia (fluid balance)

It has been suggested to limit sodium intake for its effects on both fluid balance and blood pressure.^{24, 25} In advanced CKD, the kidney is unable to adapt to large changes in sodium intake. Intake >3–4 g/day can lead to hypertension and volume overload, whereas intake of <1 g/day can lead to volume depletion and hypotension. A goal of 2 g/day of sodium is reasonable for most patients. A daily intake of 2 L of fluid maintains water balance.²⁵ But overzealous salt restriction or diuretic use can lead to extracellular fluid volume depletion and precipitate a further decline in GFR. Patients with salt-losing nephropathy may require a sodium-rich diet or salt supplementation.^{6, 24}

Hyperkalemia

Hyperkalemia often responds to dietary restriction of potassium, avoidance of potassium supplements as well as potassium-retaining medications [especially spironolactone, angiotensin-converting enzyme (ACE) inhibitors/angiotensin-receptor blockers (ARBs)], or the use of kaliuretic diuretics such as metolazone. Metolazone promotes urinary potassium excretion, while a potassium-binding resin, such as calcium resonium or sodium polystyrene, promotes potassium loss through the gastro-intestinal tract and may reduce the incidence of hyperkalemia in CKD patients.²⁵ Acute hyperkalemia: calcium chloride or gluconate intravenously, insulin administration with glucose intravenously, bicarbonate intravenously, and potassium-binding resin orally or per rectum, cardiac monitoring.²⁵ Intractable hyperkalemia is an indi-cation (although uncommon) to consider institution of dialysis in a CKD patient.²⁵ Chronic hyperkalemia:

dietary potassium restriction and sodium polystyrene, 15–30 g once daily orally in juice or sorbitol.²⁵

Table 3: Complications of CKD by stage and GFR

Stage	GFR	Description		Complications		
1	##	Kidney damage with		Patient at risk of CKD		
		normal or	GFR			
2	18>	Kidney damage with		Hypertension		
		mild	GFR			
3	10–12	Moderate	GFR	Mild	in serum creatinine, mild	in serum
				calcium, onset of anemia		
4	5–10	Severe	GFR	Anorexia, moderate anemia, hyperkalemia		
				hyperphosphatemia, dyslipidemia, neuropathy,		
				left ventricular hypertrophy		
5	<1	Kidney failure		Uremic symptoms, severe anemia,		
				malnutrition, secondary hyperparathyroidism,		
				metabolic acidosis		

Advanced chronic kidney disease, kidney failure and end stage renal disease

Stages 4 and 5 are sometimes called advanced CKD.² The NKF guidelines define kidney failure as either 1) GFR less than 15 mL/min per 1.73 m², which is accompanied in most cases by signs and symptoms of uremia, or 2) a need to start RRT.⁷ The dispiriting term ESRD represents a stage of CKD where the accumulation of toxins, fluid, and electrolytes normally.

Complications

The complications of CKD tend to occur at comparatively inevitable stages of sickness as noted in Table 3. Stages one and a pair of CKD are sometimes not related to any symptoms arising from the decrement in GFR. However, there is also symptoms from the underlying urinary organ sickness itself, like puffiness in patients with syndrome or signs of cardiovascular disease secondary to some sorts of Bright's disease. If the decline in GFR progresses to stages 3 and 4, clinical and laboratory complications of CKD become more prominent. Virtually all organ systems are affected, but the most evident complications include anemia, decreasing appetite with progressive malnutrition, abnormalities in calcium, phosphorus and mineral-regulating hormones, such as 1,25(OH)₂D₃ (calcitriol) and parathyroid hormone (PTH), and abnormalities in sodium, potassium, water and acid-base homeostasis.⁶

Acidemia

It has been suggested that in presence of acidemia oral alkali therapy can slow CKD progression.²⁵ The plasma bicarbonate should be maintained >22 mmol/L by giving sodium bicarbonate supplements (starting dose of 1 g 8-hourly, increasing as required).^{6,24} It must be remembered that the increased sodium intake may induce hypertension or edema.²⁴

Hypertension

Hypertension is commonly associated with CKD in more than 75% of patients.²⁶ Various target blood pressures have been suggested, for example, 130/80 Hg or less for CKD alone, and lowered to 125/75 mm Hg for those with proteinuria >1g/day [^{6,7, 24, 27, 28}]. In CKD if hypertension is

associated with significant proteinuria (0.5 g/24 h or more), ACE inhibitors/ ARBs should be offered because of their unique ability to decrease proteinuria preferentially by lowering intra-glomerular pressure.^{2,27} In hyper-tension with urinary protein excretion <0.5 g/24 h, ACE inhibitors, ARBs, long-acting calcium channel blockers, thiazide diuretics or beta-blockers can be used.² When ACE inhibitor/ARB therapy is started, some patients with CKD may have an initial decrease in GFR, a mild increase in the serum creatinine concentration, and a mild increase in the potassium level.²⁹ Therefore, in people with CKD, serum creatinine and potassium levels should be measured and the GFR should be estimated before starting ACE inhibitor/ARB therapy. These measurements should be repeated between 7th and 14th day after starting ACE inhibitor/ARB therapy and after each dose increase.² ACE inhibitor/ARB therapy should not normally be started if the pretreatment serum potassium concentration is significantly above the normal reference range. Often the concomitant use of a kaliuretic diuretic can improve potassium excretion in addition to improving blood pressure control.^{2, 6}

Proteinuria

Increased intraglomerular filtration pressures develop as a response to loss of nephron number from CKD as well as from systemic hypertension. Control of systemic and glomerular hypertension is important in slowing the progression of CKD. ACE inhibitors and ARBs inhibit the angiotensin-induced vasoconstriction of the efferent arterioles of the glomerular microcirculation. This inhibition leads to a reduction in both intraglomerular filtration pressure and proteinuria. ACE inhibitors or ARBs should be offered to non-diabetic people with CKD with urinary protein excretion 0.5 g/24 h or more.^{2, 27.}

Anemia

Erythropoiesis-stimulating agents (ESAs) are FDA-approved in CKD for a goal hemoglobin (Hb) of 10–11 g/dL if no other treatable causes for anemia are present.²⁵ Other various target Hb suggested by different clinical guidelines are 10–12 g/dL^{2,24}, 11–12 g/dL^{7,27} and 10–11.5 g/dL.⁶ ESAs (e.g., recombinant erythropoietin [epoetin] and darbepoetin) are effective in correcting anemia of CKD. The

starting dose of epoetin is 50 units/kg (3000–4000 units/dose) once or twice a week. Darbepoetin is started at 0.45 mcg/kg and can be administered every 2–4 weeks.²⁵ ESAs are less effective in the presence of iron deficiency, active inflammation, malignancy, intercurrent illness, or in patients with aluminium overload, which may occur in dialysis.²⁴ The use of ESAs for the treatment of anemia in patients with CKD is associated with potential adverse outcomes, including increased blood pressure, thrombotic complications (like stroke) and possibly other CVD events.^{25,27} Iron therapy is an important component of anemia management. Iron supplementation is usually essential to ensure an optimal response to ESAs in patients with CKD because the demand for iron by the marrow frequently exceeds the amount of iron that is immediately available for erythropoiesis (measured by percent transferrin saturation), as well as the amount in iron stores (measured by serum ferritin). In CKD, a serum ferritin <100 ng/mL or iron saturation <20% is suggestive of iron deficiency.²⁵ Iron stores should be repleted prior to the initiation of ESAs. For the CKD patient not yet on dialysis, oral iron supplementation should be attempted. If there is GI intolerance, the patient may have to undergo intravenous (IV) iron infusion.⁶ People receiving ESA maintenance therapy should be given iron supplements to keep their serum ferritin levels between 200 and 500 ng/mL.² Iron therapy should probably be withheld if the serum ferritin is >500 ng/mL, even if the iron saturation is <20%.²⁵ Oral therapy with ferrous sulfate, gluconate, or fumarate, 325 mg once to three times daily, is the initial therapy in CKD.²⁵ For patients on hemodialysis, IV iron can be administered during dialysis. In addition to iron, an adequate supply of other major substrates and cofactors for red cell production must be ensured, including vitamin B12 and folate.²⁵ Red cell transfusion to treat anemia in CKD becomes necessary sometimes. In people with anemia of CKD in whom kidney transplant is a treatment option, red cell transfusions should be avoided if possible, to minimize the risk of allo-immunization related to their use.² It has been suggested by investigators in one study that the decision to transfuse a CKD patient with non-acute anemia should not be based on any arbitrary Hb threshold only, but should be determined also by the occurrence of symptoms caused by anemia. In particular, they suggest considering transfusion at Hb values ## g/dL or at Hb values ## g/dL in postoperative surgical patients.³⁰

4. Discussion

In CRF, hypocalcemia results from inadequate production of 1, 25 - dihydroxycholecalciferol (calcitriol) by the kidneys needed for calcium retention and resorption. Hyperphosphatemia is a consequence of reduced urinary phosphate excretion due to declining GFR. Phosphorus complexes with calcium, deposits in soft tissues and causes soft tissue calcification. Hypocalcemia, decreased calcitriol and hyperphosphatemia stimulates the parathyroid glands for secretion of excessive parathyroid hormone (PTH) resulting in secondary hyperparathyroidism. This typical pattern of hyperphosphatemia, hypocalcemia, hypovitaminosis D, and secondary hyperparathyroidism is seen as early as CKD stage 3.²⁵ Excessive PTH helps to maintain calcium levels mainly by bone resorption and causes osteitis fibrosa and other disturbances like osteomalacia, osteoporosis and osteosclerosis.²⁴ These

disturbances have traditionally been termed renal osteodystrophy.³¹ Disturbances in mineral metabolism also results in vascular calcification due to deposition of calcium phosphate complex, resulting in vascular rigidity and occlusion which may be partly responsible for the accelerated CVD and excess mortality seen in the CKD population. As the disorder is not confined to the skeleton, in 2005, the Kidney Disease: Improving Global Outcomes (KDIGO) clinical practice proposed replacing the term renal osteodystrophy by CKD-MBD.³² CKD-MBD is manifested by an abnormality of any one or a combination of the following disorders-- laboratory-abnormalities of calcium, phosphorus, PTH, or vitamin D metabolism; bone-changes in bone turnover, mineralization, volume, or strength and calcification in the soft-tissue and the arterial vessels.^{31,32} Control of hyperphosphatemia is the first step in the treatment of CKD-MBD.²⁵ This involves dietary phosphorus restriction initially, followed by the administration of oral phosphorus binders if targets are not achieved. Examples of foods with high phosphate content are milk, cheese, eggs, nuts, peas, beans, colas, and dairy products.²⁴ Oral phosphorus binders, such as calcium carbonate/gluconate (500 mg/tablet, maximum 3 gm daily) or calcium acetate (667 mg/capsule, maximum 1500 mg daily) block absorption of dietary phosphorus in the gut and are given thrice daily with meals. These should be titrated to a serum phosphorus of <4.5 mg/dL in stage 3–4 CKD and <5.5 mg/dL in ESRD. A major side-effect of calcium-based phosphate binders is hypercalcemia. Therefore doses should be decreased or stopped if serum calcium rises above 10 mg/dL.²⁵ Phosphorus-binding agents that do not contain calcium are aluminium hydroxide, sevelamer and lanthanum. Sevelamer, 800–3200 mg and lanthanum carbonate, 500–1000 mg are given at the beginning of meals and may be combined with calcium- containing binders.^{6, 25} Aluminium exposure should be avoided. High levels of aluminium may be present in impure dialysate water.²⁵ Aluminium hydroxide is a highly effective phosphorus binder, but can cause osteomalacia and neurologic complications when used long-term. It can be used in the acute setting for serum phosphorus >7 mg/dL or for short periods (e.g., 3 weeks) in CKD patients; however, long-term use should be avoided.²⁵ Secondary hyperparathyroidism can be prevented by restricting dietary phosphate intake, using a calcium-based phosphate binder, and administering calcitriol.³³ Best practice is to maintain normal calcium, phosphate, and 1, 25-dihydroxycholecalciferol. Once serum phosphorus levels are controlled, active vitamin D (calcitriol) or active vitamin D analogues are recommended to treat secondary hyperparathyroidism (if PTH more than twice normal) or hypocalcaemia in stage 3–5 CKD.^{24,25} Typical calcitriol dosing is 0.25 or 0.5 mcg orally daily or every other day.²⁵ Serum 25-OH vitamin D levels should be measured and brought to normal by vitamin D supplementation.³⁴ When vitamin D supplementation is indicated in people with CKD, certain clinical guidelines recommend cholecalciferol or ergocalciferol to people with stage 1, 2, or 3 CKD and calcitriol or 1-alpha-hydroxycholecalciferol (alfacalcidol) to people with stage 4 or 5 CKD.³⁵ The routine measurement of serum calcium, phosphate, and PTH levels in people with stage 1, 2, 3 CKD is not recommended, but needs measurement in people with stage 4 or 5 CKD.³⁵ Calcitriol therapy may result in hypercalcemia and/or hyperphosphatemia through increased

GI absorption of these minerals. The dose of calcitriol should be decreased or stopped if hypercalcemia or hyperphosphatemia occurs.² Calcitriol exerts a direct suppressive effect on PTH secretion and also indirectly suppresses PTH secretion by raising the concentration of ionized calcium. Certain analogues of calcitriol (e.g., paricalcitol) are available that suppress PTH secretion with less attendant hypercalcemia.⁶ Calcimimetic agents (e.g., cinacalcet) which enhance the sensitivity of the parathyroid cell to the suppressive effect of calcium have been developed recently. This class of drug produces a dose-dependent reduction in PTH production and plasma calcium concentration.^{6, 35} Cinacalcet, 30–90 mg orally once a day, can be used if elevated serum phosphorus or calcium levels prohibit the use of vitamin D analogues.²⁵ In 2009 KDIGO clinical practice released guidelines on the management of CKD-MBD which suggests that, if the PTH levels fall below two times the upper limit of normal for the assay, calcitriol or calcimimetics be reduced or stopped.^{6, 35}

Tertiary hyperparathyroidism: Even with appropriate medical therapy, some patients continue to have refractory hyperparathyroidism because of parathyroid gland hyperplasia. In patients with severe hyperparathyroidism who fail to respond to medical/pharmacological therapy, the KDIGO guideline suggests parathyroidectomy.³⁵

Dyslipidemia

Dyslipidemia is a common complication of progressive CKD and a primary risk factor for CVD. The goals are an LDL cholesterol level below 100 mg per dL (2.60 mmol per L) and a triglyceride level below 200 mg per dL (2.26 mmol per L).³⁶ Statins should be offered to people with CKD for the secondary prevention of CVD irrespective of baseline lipid values.² Fibrates are known to decrease triglyceride levels, but they may increase the risk for rhabdomyolysis in patients with CKD.²⁷

Antiplatelet drugs

Antiplatelet drugs should be offered to people with CKD for the secondary prevention of CVD.²

Hyperuricemia

There is also theoretic value in lowering uric acid levels in CKD patients with concomitant hyperuricemia, but clinical data regarding benefit are still lacking.²⁵

Nutrition

Counseling on adequate and appropriate nutrition should be provided at each office visit. Dietary advice should be offered to people with progressive CKD concerning protein, calorie, salt, potassium and phosphate intake, when indicated.² Protein intake: The effect of dietary protein restriction on kidney disease is the subject of debate. There is a lack of convincing evidence that the long-term restriction of protein intake (<0.70 g/kg/day) delays the progression of CKD.^{6, 27} Nonetheless, restriction of dietary protein intake has been recommended for CKD patients. For non-diabetic and diabetic patients with CKD stage 4, two systematic reviews and one meta-analysis suggested that, in comparison to other treatments, there was, at most, a modest benefit associated with restricting protein leading to a delay in CKD progression.^{37,38} High protein intakes are associated with high phosphate intakes as foods that contain protein also tend to contain phosphate. It would appear

prudent to avoid high protein intakes in stage 4 CKD patients when hyperphosphatemia is prevalent. The ADA recommends reduction of protein intake to 0.8–1.0 g/kg body wt/day in individuals with diabetes and the earlier stages of CKD and to 0.8 g/kg body wt/day in the later stages of CKD that may improve renal function.³⁹ Current recommendations from the NKF K/DOQI and other studies suggest a protein intake of 0.8 to 1.0 g/kg/day in patients with CKD.^{7,40–42} It is further advised that at least 50% of the protein intake be of high biologic value.⁶ Benefits of protein restriction in slowing decline of GFR must be weighed against the risk of malnutrition and cachexia upon the initiation of dialysis.²⁵ Energy intake: An adequate nutrition with daily caloric intake of 30 to 35 kcal per kg/day in patients with CKD has been recommended.^{6,7} Salt and water restriction: A goal of 2 g/day of sodium intake is reasonable for nondialysis patient approaching ESRD; a daily intake of 2 L of fluid maintains water balance.²⁵ Patients with salt-losing nephropathy may require a sodium-rich diet or salt supplementation.⁶ Potassium restriction: A goal of <50–60 mEq/day intake of potassium is adequate. Restriction is needed once the GFR has fallen below 10–20 mL/min/1.73 m², or earlier if the patient is hyperkalemic.²⁵ Phosphorus restriction: Phosphorus needs to be restricted to 800–1000 mg/day. Foods rich in phosphorus such as cola beverages, eggs, dairy products, nuts, beans, and meat should be limited, although care must be taken to avoid protein malnutrition.²⁵ Phosphorus-containing medicines, particularly cathartics should be avoided.²⁵

Magnesium-containing medications, such as laxatives and antacids are contraindicated. Many drugs are excreted by the kidney; dosages should be adjusted for GFR.²⁵

Lifestyle advice

Most patients with CKD will die of events related to CVD before ESRD develops.⁴³ Therefore, an important focus of care for patients with CKD includes management of cardiovascular risk factors. People with CKD should be encouraged at each office visit to take regular exercise, achieve a healthy weight and stop smoking completely. Smoking is a strong risk factor for cardiovascular mortality in patients with CKD.⁴⁴ It is also strongly associated with the progression of nephropathy.⁴⁴ A target weight of BMI <23 should be ideal.²

Treatment of underlying cause for preventing progression of CKD

Treatment of the underlying cause of CKD is vital. These include optimized glucose control in diabetes mellitus, immunomodulatory agents for glomerulonephritis, and emerging specific therapies to retard cytogenesis in polycystic kidney disease.^{6, 25}

Diabetes

Control of diabetes should be aggressive in early CKD; excellent glycemic control reduces the risk of kidney disease and its progression.²⁵ The American Diabetes Association (ADA) and Canadian guidelines recommend that plasma values for preprandial glucose be kept in the 5.0–7.2 mmol/L (90–130 mg/dL) range and hemoglobin A1C should be <7% to prevent progression of CKD in patients with DM.^{27,45} The half-life of insulin is prolonged in CKD due to reduced tubular metabolism of insulin;

patients on insulin therapy may need progressive reduction in dose as their renal function worsens. Risk of hypoglycemia increases in advanced CKD, and glycemic targets may need to be relaxed to avoid this dangerous complication.²⁴ Many hypoglycemic agents such as metformin require dose reduction in renal failure, and some are contraindicated when the GFR is <50 mL/min/1.73 m².²⁵ Metformin can cause lactic acidosis; and the thiazolidinediones (e.g., pioglitazone) may increase renal salt and water absorption and aggravate volume-overload states, and contribute to adverse cardiovascular events.⁶ ACE inhibitors or ARBs should be used in individuals with microalbuminuria as well as with macroalbuminuria irrespective of presence of hypertension in diabetes.⁹ ACE inhibitors and ARBs are effective at reducing proteinuria, slowing the decline in GFR and retarding the progression of diabetic kidney disease.² The definition of hypertension in DM or diabetic kidney disease (DKD) is blood pressure of 130/80 Hg or greater.⁴⁶ In patients with diabetes having kidney disease and hypertension, treatment with ACE inhibitors or ARBs has been recommended by various guidelines to prevent progression of CKD.^{2, 8-13}

Initiation of renal replacement therapy (RRT)

Although the overall objectives of the guidelines and recommendations are to delay progression of both CKD and its complications, kidney function may continue to deteriorate in a proportion of patients. Ultimately, patients may develop uremia and kidney failure and the need of RRT (either dialysis or transplantation) to extend their life. Patients with kidney failure should be evaluated for kidney transplantation. A kidney transplant from a live donor should be promoted as the first choice for eligible patients who require RRT. Outcomes with deceased donor kidneys are also significantly better than with dialysis. For patients with progressive CKD with plans for hemodialysis, making plans for vascular access is an important component of their care in preparation for ESRD. Acute uremia or uremia resulting from progressive disease is an indication for immediate dialysis.

5. Conclusion

The incidence of CKD is increasing worldwide and unfortunately, CKD is often undetected and undertreated because of its insidious onset, variable progression, and length of time to overt kidney failure. Its natural course is an eventual decline in the GFR, leading to kidney failure and the need for RRT. The goal of therapy is to detect CKD early to delay its progression, thereby reducing associated complications and adverse outcomes of kidney disease. Patients with one or more risk factors should have their GFR estimated and be assessed for persistent albuminuria. Evidence supports that achieving optimal glucose control, blood pressure, and reduction in albuminuria slows the progression of CKD.^{7, 10, 28, 55-59} ACE inhibitors are more effective than other antihypertensive drugs in preventing the progression of kidney disease in diabetic and nondiabetic patients. Other interventions studied to reduce progression include correction of anemia, treatment of CKD-MBD, dietary protein restriction, and lipid-lowering therapy.

6. Recommendations

Early detection and management of CKD can prevent kidney disease progression and decrease its complications. All patients should be assessed to determine their susceptibility to or development of CKD. The clinical evaluation should include analysis for risk factors, markers of CKD (e.g. assessment for persistent albuminuria), GFR estimation, and a treatment plan to slow the progression. Key clinical recommendations for treatment, prevention and delaying the progression of CKD and its complications are: According to the evidence, ACE inhibitors/ARBs have been shown to reduce proteinuria and the occurrence of kidney failure.^{2,29,47-50} Evidence suggests that ACE inhibitors are more effective than other antihypertensive drugs in preventing the progression of kidney disease in diabetic and nondiabetic patients.⁵¹⁻⁵⁴ The shishu hospital seventh report of the Joint Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure recommends a target blood pressure of less than 130/80 mm Hg in patients with chronic kidney disease.²⁸ To prevent progression of nephropathy in patients with DM, the ADA recommends glycemic control, with the goal being a HbA1C <7 percent.⁴⁵ The most recent guidelines from the NKF recommend treating dyslipidemia aggressively in patients with CKD. The goals are a LDL cholesterol level below 100 mg per dL (2.60 mmol/L) and a triglyceride level below 200 mg per dL (2.26 mmol/L).³⁶ Anemia in CKD responds to ESAs. Iron therapy is an important component of anemia management in CKD. The target goal of Hb is 10–11 g/dL.²⁵ In CKD patients, where kidney transplant is a treatment option, red cell transfusions should be avoided if possible.² Epidemiologic studies in humans show an association between elevated phosphorus levels and increased risk of cardiovascular mortality in early CKD through ESRD.²⁵ Control of hyperphosphatemia is the first step in the treatment of CKD-MBD.²⁵ While using calcium-based phosphorus binders, hypercalcemia must be avoided to prevent aggravation of soft tissue and vascular calcification leading to increased cardiovascular morbidity and mortality. Protein restriction may be effective in slowing the progression of CKD, especially proteinuric and diabetic renal diseases. Benefits of protein restriction in slowing decline of GFR must be weighed against the risk of malnutrition and cachexia upon the initiation of dialysis.²⁵

Conflict of interest: The Author has no conflict of interest of the study.

7. References

1. Excerpts from the United States renal data system's 2000 annual data report: atlas of end-stage renal disease in the United States. *Am J Kidney Dis.* 2000; 36(6Suppl 2):S1-S137?
2. NICE clinical guidelines. Chronic kidney disease. Early identification and management of chronic kidney disease in adults in primary and secondary care CG73. September 2008. Available at: <http://publications.nice.org.uk/chronic-kidney-disease-cg73>. Accessed July, 2013.
3. Rashid HU. Bangladesh renal registry report. 1986-1996. *Bang Renal J.* 2002; 21(1):25-28.

4. Sakhujia V, Sud K. End stage renal disease in India and Pakistan: burden of disease and management issues. *Kidney Int Suppl.* 2002; 83:S115-S118.
5. Ahmed ST, Rahim MA, Ali MZ, and Iqbal MM. Prevalence of primary renal disease among patients on maintenance hemodialysis: a hospital based study. *KYAMC Journal.* 2012; 2(2):182-186.
6. Bargman JM, Skorecki K. Chronic kidney disease. In: Longo DL, Fauci AS, Kasper DL, Hauser SL, Jameson JL, Loscalzo J (eds). *Harrison's principles of internal medicine.* 18th edn. New York: McGraw-Hill, 2012, 2308-2321.
7. National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis.* 2002; 39(2 Suppl 1):S1-S266?
8. UK CKD guidelines consultation draft 9th December 2005. Available at: http://www.kidney.org.uk/main/guidelines/UK_CKD_guidelines_consultation_draft.pdf. Accessed May, 2013.
9. International Diabetes Federation, 2012. Clinical guidelines task force global guideline for type 2 diabetes. Available at: <http://www.societate-diabet.ro/pdf/global-guideline-for-type-2-diabetes-IDF-2012.pdf>. Accessed June, 2013.
10. American Diabetes Association. Position statement: nephropathy in diabetes. *Diabetes Care.* 2004; 27:S79-S83.
11. Canadian Diabetes Association Clinical Practice Guidelines Expert Committee. *Canadian Diabetes Association 2008 clinical practice guidelines for the prevention and management of diabetes in Canada.* *Can J Diabetes.* 2008; 32:S95-S98.
13. Scottish Intercollegiate Guidelines Network. Management of diabetes. A national clinical guideline. March 2010. Available at: <http://www.sign.ac.uk/pdf/sign116.pdf>. Accessed March, 2013.
14. Chadban S, Howell M, Twigg S, Thomas M, Jerums G, Alan C, *et al.* National evidence based guideline for diagnosis, prevention and management of chronic kidney disease in type 2 diabetes. Canberra: Diabetes Australia and the NHMRC; 2009. Available at: http://www.nhmrc.gov.au/_files_nhmrc/publications/attachments/di18-diabetes-kidney-disease.pdf. Accessed April, 2013.
15. Snively SC, Gutierrez C. Chronic kidney disease: prevention and treatment of common complications. *Am Fam Physician* Cumming A, Payne S. The renal system. In: Douglas G. 2004; 70(10):1921-1928.
16. Nicol F, Robertson C (eds). *Macleod's clinical examination.* 12th edn. Edinburgh: Churchill Livingstone Elsevier, 2009, 223-224.
17. Levey AS, Coresh J, Balk E, Kausz AT, Levin A, Steffes MW *et al.* National Kidney Foundation practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Ann Intern Med.* 2003, 137-147.
18. Levey AS, Bosch JP, Lewis JB, Greene T, Rogers N, Roth D. A more accurate method to estimate glomerular filtration rate from serum creatinine: a new prediction equation. Modification of diet in renal disease study group. *Ann Intern Med.* 1999; 130:461-470.
19. Davies DF, Shock NW. Age changes in glomerular filtration rate, effective renal plasma flow, and tubular excretory capacity in adult males. *J Clin Invest.* 1950, 496-507.
20. Lindeman RD, Tobin J, Shock NW. Longitudinal studies on the rate of decline in renal function with age. *J Am Geriatr Soc.* 1985; 33:278-285.
21. Rowe JW, Andres R, Tobin JD, Norris AH, Shock NW. The effect of age on creatinine clearance in men: a cross-sectional and longitudinal study. *J Gerontol.* 1976, 155-163.
22. Cockcroft DW, Gault MH. Prediction of creatinine clearance from serum creatinine. *Nephron.* 1976; 16:31-41.
23. Mendelssohn DC. Policy statement — principles of end-stage renal disease care. *Annals RCPSC.* 1997; 30(5):271-273.
24. Parmar M. Chronic renal disease. *BMJ.* 2002; 325:85-90, 54.
25. Goddard J, Turner AN, Stewart LH. Kidney and urinary tract disease. In: Colledge NR, Walker BR, Ralston SH (eds). *Davidson's principle and practice of medicine.* 21st edn. Edinburgh: Churchill Livingstone Elsevier, 2010, 480-496.
27. Watnick S, Dirks T. Kidney disease. In: Papadakis MA, McPhee SJ, Rabow MW (eds). *Current medical diagnosis & treatment 2013.* New York: McGraw-Hill, 2013: Chapter 22. Available: at <http://www.accessmedicine.com/content.aspx?aID=11374>. Accessed August, 2013.
28. Culleton B. Introduction to the Canadian clinical practice guidelines. *J Am Soc Nephrol.* 2006; 17:S1-S3.
29. Levin A, Hemmelgarn B, Culleton B, Tobe S, McFarlane P, Ruzicka M *et al.* Guidelines for the management of chronic kidney disease. *CMAJ.* 2008; 179(11):1154-1162.
30. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr *et al.* The seventh report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure. *JAMA.* 2003; 289:2460-2472.
31. Levey AS. Clinical practice. Nondiabetic kidney disease. *N Engl J Med.* 2002; 347:1505-1511.
32. Drüeke TB, Locatelli F, Clyne N, Eckardt KU, Macdougall IC, Tsakiris D *et al.* CREATE investigators. Normalization of hemoglobin level in patients with chronic kidney disease and anemia. *N Engl J Med.* 2006; 355:2071-2084.
33. Moe SM, Drüeke T, Lameire N, Eknoyan G. Chronic kidney disease-mineral-bone disorder: a new paradigm. *Adv Chronic Kidney Dis.* 2007; 14(1):3-12.
34. Suchowierska E, Mysliwiec M. Mineral and bone disturbances associated with chronic kidney disease [Review]. *Pol Merkur Lekarski.* 2010; 28(164):138-143. PMID: 20369744.
35. Ho LT, Sprague SM. Renal osteodystrophy in chronic renal failure. *Semin Nephrol.* 2002; 22:488-493.
36. Slatopolsky E, Berkoben M, Kelber J, Brown A, Delmez J. Effects of calcitriol and non-calcemic vitamin D analogs on secondary hyperparathyroidism. *Kidney Int Suppl.* 1992; 38:S43-S49.
37. Moe SM, Drüeke TB, Block GA, Cannata-Andía JB, Elder GJ, Fukagawa M *et al.* KDIGO clinical practice guideline for the diagnosis, evaluation, prevention, and

- treatment of chronic kidney disease-mineral and bone disorder (CKD-MBD). *Kidney Int Suppl.* 2009; 113:S1-S130.
38. National Kidney Foundation. Kidney Disease Outcomes Quality Initiative. K/DOQI clinical practice guidelines for managing dyslipidemias in chronic kidney disease, 2003. Available at: http://www.kidney.org/professionals/kdoqi/guidelines_lipids/index.htm. Accessed July, 2013.
 39. Fouque D, Wang P, Laville M, Boissel JP. Low protein diets for chronic kidney disease in nondiabetic adults (Cochrane review). In: *The Cochrane Database Syst Rev*, 2009, Issue 3. DOI: 10.1002/14651858.CD001892.pub3.
 40. Robertson L, Waugh N, Robertson A. Protein restriction for diabetic renal disease (Cochrane review). In: *The Cochrane Database Syst Rev*, 2007, 4. CD002181.
 41. American Diabetes Association. Standards of medical care in diabetes. *Diabetes Care.* 2013; 36(Suppl 1):S11-S66.
 42. Klahr S, Levey AS, Beck GJ, Caggiula AW, Hunsicker L, Kusek JW *et al.* The effects of dietary protein restriction and blood-pressure control on the progression of chronic renal disease. Modification of Diet in Renal Disease Study Group. *N Engl J Med.* 1994; 330:877-884.
 43. Fouque D, Laville M, Boissel JP. Low protein diets for chronic kidney disease in nondiabetic adults. *Cochrane Database Syst Rev.* 2006; 2:CD001892.
 44. Kasiske BL, Lakatua JD, Ma JZ, Louis TA. A meta-analysis of the effects of dietary protein restriction on the rate of decline in renal function. *Am J Kidney Dis.* 1998; 31:954-961.
 45. Keith DS, Nichols GA, Gullion CM, Brown JB, Smith DH. Longitudinal follow-up and outcomes among a population with chronic kidney disease in a large managed care organization. *Arch Intern Med.* 2004; 164:659-663.
 46. Orth SR, Stockmann A, Conradt C, Ritz E, Ferro M, Kreusser W *et al.* Smoking as a risk factor for end-stage renal failure in men with primary renal disease. *Kidney Int.* 1998; 54:926-931.
 47. American Diabetes Association. Standards of medical care for patients with diabetes mellitus. *Diabetes Care.* 2003; 26(Suppl 1):S33-S50.
 48. National Kidney Foundation. K/DOQI Clinical practice guidelines and clinical practice recommendations for diabetes and chronic kidney disease. *Am J Kidney Dis.* 2007; 49(2 Suppl 2):S1-S179.
 49. Mogensen CE, Neldam S, Tikkanen I, Oren S, Viskoper R, Watts RW *et al.* Randomised controlled trial of dual blockade of the renin-angiotensin system in patients with hypertension, microalbuminuria, and non-insulin dependent diabetes: the Candesartan and Lisinopril Microalbuminuria (CALM) study. *BMJ.* 2000; 321:1440-1444.
 50. Siebenhofer A, Plank J, Horvath K, Berghold A, Sutton AJ, Sommer R *et al.* Angiotensin receptor blockers as anti-hypertensive treatment for patients with diabetes mellitus; meta-analysis of controlled double-blind randomized trials. *Diabet Med.* 2004; 21:18-25.
 51. Praga M, Andrade CF, Luno J, Arias M, Poveda R, Mora J *et al.* Antiproteinuric efficacy of losartan in comparison with amlodipine in non-diabetic proteinuric renal diseases: a double-blind, randomized clinical trial. *Nephrol Dial Transplant.* 2003; 18:1806-1813.
 52. Brenner BM, Cooper M, de Zeeuw D, Keane WF, Mitch WE, Parving HH *et al.* Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N Engl J Med.* 2001; 345:861-869.
 53. Kasiske BL, Kalil RS, Ma JZ, Liao M, Keane WF. Effect of antihypertensive therapy on the kidney in patients with diabetes: a meta-regression analysis. *Ann Intern Med.* 1993; 118:129-138.
 54. Wright JT, Bakris G, Greene T, Agodoa LY, Appel LJ, Charleston J *et al.* Effect of blood pressure lowering and antihypertensive drug class on progression of hypertensive kidney disease: results from the AASK trial. *JAMA.* 2002; 288:2421-2431.
 55. Randomised placebo-controlled trial of effect of ramipril on decline in glomerular filtration rate and risk of terminal renal failure in proteinuric, non-diabetic nephropathy. The GISEN Group (Gruppo Italiano di Studi Epidemiologici in Nefrologia). *Lancet.* 1997; 349:1857-1863.
 56. Ruggenenti P, Perna A, Gherardi G, Gaspari F, Benini R, Remuzzi G, *et al.* Renal function and requirement for dialysis in chronic nephropathy patients on long-term ramipril: REIN follow-up trial. Gruppo Italiano di Studi Epidemiologici in Nefrologia (GISEN). *Ramipril Efficacy in Nephropathy.* *Lancet.* 1998; 352:1252-1256.
 57. K/DOQI clinical practice guidelines on hypertension and antihypertensive agents in chronic kidney disease. *Am J Kidney Dis.* 2004; 39:S11-S68.
 58. Diabetes Control and Complications Trial (DCCT) Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med.* 1993; 329:977-986.
 59. UK Prospective Diabetes Study Group. Intensive blood glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet.* 1998; 352:837-853.
 60. American Diabetes Association. Position statement: hypertension management in adults with diabetes. *Diabetes Care.* 2004; 27:S65-S67.
 61. Manley HJ. Role of angiotensin-converting-enzyme inhibition in patients with renal disease. *Am J Health-Syst Pharm.* 2000; 57(Suppl 1):S12-S18.