



Experiences of acute myocardial infarction survivors: A qualitative literature review

Puneet Kaur¹, Dr. DK Seth², Dr. Sukhpal Kaur³

¹ Lecturer KCON Kasturba Hospital Delhi, New Delhi, India

² Principal Consultant, MCD Delhi, New Delhi, India

³ Associate Professor, PGIMER Chandigarh, India

Abstract

Background Aims: Cardiovascular diseases are among one the leading causes of death worldwide. The advancements in technology have improved survival rates in acute myocardial infarction patients. Survivors of acute myocardial infarction patients have to face many challenges to adapt to new situations. In-depth understanding of their lived experiences is pivotal for designing tailored made educational programme for their prevention and rehabilitation. There are limited researches available and inadequate attention has been paid to this issue. Therefore, this paper will evaluate personal lived experiences and perspectives of acute myocardial infarction survivors.

Material and Methods: Qualitative Literature Review from DELNET, MEDLINE, CINHAI and EBESCO electronic databases. Nineteen articles, 2002-2018, met the inclusion criteria. To identify themes constant comparative method was used.

Results: Six key themes were identified: knowledge of causes (risk factors) of the disease, responses led by the event, adopting lifestyle changes, family and social support, future concerns and insecurities and expression of needs. Many survivors did not have adequate knowledge pertaining to causes and risk factors, warning signs and symptoms of acute myocardial infarction. Participants found changing one's lifestyle is difficult and require constant support from family and health care providers.

Conclusion: Community awareness campaigns are required to be increased by health care providers. Patients should be educated about the risk factors, signs, symptoms and appropriate changes in life style during their routine checkup visits. Acute myocardial survivors need time-sensitive education especially during and after their hospitalization. Patients need constant support from family and health care providers' to combat challenges during their recovery process. Health care providers can utilize this information to develop teaching programmes and to assist in care of acute myocardial infarction survivors.

Keywords: myocardial infarction, lifestyle changes, literature review, survivors, teaching programme

Introduction

Cardiovascular diseases (CVDs), especially coronary heart disease (CHD) is one of the leading causes of morbidity and mortality throughout the world [1]. It has assumed epidemic proportions worldwide. CHD global fatality was estimated to be 17.5 million/year, 31% of deaths - 75% in low- and middle-income countries [2]. According to WHO 7.4 million deaths were due to coronary heart diseases in 2015. In contrast to developed countries mortality from CVD is increasing in developing countries. Eighty-two percent of deaths in low and middle-income countries are accountable for CVD. It is estimated that 23.6 million people will die from CVDs by 2030 [1].

The international number one killer disease (CVD) is largely a product of unhealthy lifestyle transformations in terms of urbanization, globalization and industrialization [3]. AMI is one of the manifestations of CHD that effects large number of people. In India in all age specific groups, prevalence of MI is higher in men than in women [4]. India is currently experiencing high AMI morbidity and mortality rates from AMI. In India, out of all deaths nearly 24.8% deaths are due to cardiovascular diseases as estimated by Global Burden of Disease Study (2010) [5]. The prevalence of CHD in rural India is 3%–4%, whereas it is 8%–10% in urban areas [6, 7].

By, 2020, 111% rise in deaths due to CVDs has been predicted in India [8].

Advancements in medical technology and treatment modalities have contributed towards improvement of survival rates of patients with AMI. Survivors of AMI face myriad of challenges in a way to adopt new lifestyle. Health care providers have also designed specific programs to help AMI survivors to overcome the negative consequences of the event. These programs aim to improve health outcomes and prevent other cardiac events in future.

Understanding the responses and lived experiences of AMI survivors is an essential requirement for designing and improving tailor made effective cardiac rehabilitation programs. Qualitative data provide a strong basis for understating personal and lived experiences of AMI survivors. It allows the examination of the behavior, experiences and characteristics of the study participants. Therefore, the need and purpose of this qualitative literature review is to explore AMI survivors' perspectives as their personalized experiences. This data will be useful for health care providers to develop programs to decrease incidence of AMIs and to develop effective and appropriate cardiac rehabilitation plans. Personalized patient's plan will contribute to improve the survivors' post AMI experience.

Methods

Search Strategy

Electronic databases DELNET, MEDLINE, CINHAI and EBESCO were searched for related qualitative literature, using the following keywords, qualitative research, myocardial infarction, lifestyle changes, patient’s perception, survivor’s needs. The search was restricted to articles published between 2002-2018. It was assumed that newer findings would highlight improved awareness about CVD’s, in view of the international efforts to improve CVD outcomes, with subsequent needs for support for survivors.

Inclusion Criteria

Studies meeting inclusion criteria were included: they were full texted articles; published in peer reviewed journals; qualitative research methodology articles, articles on patients’ experiences of AMI, survivors of AMI. Quantitative studies and integrative literature reviews and studies which focused on patients’ experiences on CVD in general were excluded. The flow of task during the current literature review was organized according to Polit and Beck’s task flow plan [9].

The primary search identified 80 potential articles. The abstract of each article was screened for appropriateness, which resulted in 20 articles that appeared to satisfy the inclusion criteria. However, 3 of those articles were rejected after in depth analysis. Two additional articles were included after screening the reference list of the selected articles. Finally 19 articles met the inclusion criteria.

Since, this is an integrative literature review; the authors needed no institutional review boards’ approval for the same.




Fig 1: Inclusion Scheme of Potential Articles

Data Analysis

After reviewing data of each primary source it was reduced to a single page. Following this all the primary sources were organized in a table to facilitate ordering, coding, categorizing and summarizing the data (Table-1). In the next step of data analysis, the themes from each individual source were ordered and sorted to identify common themes that reflected all the primary sources.

Strauss and Corbin’s constant comparative method was used for data comparison to assemble similar findings from every primary source under common themes [10]. This technique identified six key themes. The results were presented in a manner to help readers to understand the perspectives of patients’ about their post AMI experiences.

Table 1: Authors Name, Sample size, Qualitative Research Methodology, Data Collection Methods and Data Analysis

Author/Country	n	Gender/ Age (Mean)	Sampling Criteria	Research Methodology	Data Collection Method	Data Analysis
Kerr <i>et al.</i> (2002) Canada	7	7F 76-86yrs	Purposive	Heideggerian Phenomenology	Unstructured Interviews	Interpretive process of hermeneutic analysis
Webster <i>et al.</i> (2002)UK	35	25M/10F 65yrs	Purposive	Ethnography	Semi structured Interviews	Thematic Analysis
Bergan <i>et al.</i> (2003) Sweden	13	7M/ 6F 38-75yrs	Purposive and Theoretical	Grounded Theory	Unstructured Interviews	Grounded theory analysis technique
Jensen <i>et al.</i> (2003) Denmark	30	22M/ 8F 41-80yrs	Purposive	Not Specified	Semi structured Interviews	Content analysis
Clark (2003) UK	14	8M/6F 36-86yrs	Purposive	Not Specified	Semi structured Interviews	With specific focus on the language
Svedlund <i>et al.</i> (2004) Sweden	9	9F 47-57yrs	Purposive	Hermeneutic Phenomenology	Semi structured Interviews	Hermeneutic phenomenological mode of analysis
Hanssen <i>et al.</i> (2005) Norway	14	12M/2F 42-69 yrs	Purposive	Not Specified	Three focus groups	QSR NS software, Content analysis
Gregory <i>et al.</i> (2005) Scotland	53	35M/18W	Purposive	Not Specified	Focus group discussion	Thematic analysis
Brink <i>et al.</i> (2006) Sweden	21	10M/11F 48-83 yrs	Purposive and Theoretical	Grounded Theory	Unstructured Interviews	Grounded theory analysis technique
Condon <i>et al.</i> (2006) Ireland	10	9M/1F 38-75yrs	Purposive	Descriptive Qualitative	Semi structured Interviews	Thematic Analysis
Maclnnes (2006) UK	10	10F 72yrs	Purposive	Not Specified	Semi structured Interviews	Thematic analysis
Johansson <i>et al.</i> (2007) Sweden	15	9M/6F 41-83 yrs	Purposive	Phenomenology	Semi structured Interviews	Phenomenological method of analysis
Albarran <i>et al.</i> (2007) UK	12	12F 47-57yrs	Purposive	Not Specified	Semi structured Interviews	Thematic Analysis
Kristofferzon <i>et al.</i> (2007) Sweden	39	19M/20F 47-88yrs	Purposive	Not Specified	Semi structured Interviews	Content analysis
Moonaghi <i>et al.</i> (2014) Iran	18	12M/6W 45-80yrs	Purposive	Not Specified	Semi structured Interviews	Content analysis
Christopher <i>et al.</i> (2017) UK	10	Under 45	Purposive	Phenomenology	Semi structured Interviews	Interpretative phenomenological analysis

Sharovsky <i>et al.</i> (2017) Brazil	8	8M 48-70yrs	Purposive	Not Specified	Semi structured Interviews	Content analysis
Nicolai <i>et al.</i> (2018) Germany	21	13M/8F	Purposive	Not Specified	Semi structured Interviews	Content analysis

Results

Among the 19 qualitative studies, eight authors reported that they used qualitative tradition to conduct their studies: four studies used a phenomenological approach [11, 13, 16, 19, 20, 22, 26], two used the grounded theory approach [13, 19], one was a descriptive qualitative study [20], and one was ethnographic study [12]. Purposive sampling was used in all studies and the two grounded theory studies incorporated theoretical sampling for representation of themes [13, 19]. 10 studies included more males than females in the sample, four studies included only females population, [11, 16, 21, 23] two studies only used male participants [26, 27], two studies included more women than men in sample [19, 24]. (Table 1) Audio-tape recording was used for data recording in all the studies. 14 studies collected data through semi-structured interview [12, 14, 16, 20, 28], three through un-structured interviews [11, 13, 19] and two used the focus group technique [17, 18]. Data was analyzed manually according to traditional qualitative analysis technique.

Six key themes were identified: anticipation about causes (risk factors) of disease, responses related to the event, adopting lifestyle changes, family and social support, future concerns and insecurities and expression of needs.

Knowledge of Causes (Risk Factors) of Disease

Majority of participants stated that stress has become part of their daily living and they identified stress as the major root source of AML [14, 17]. They suffered from physical and physiological stress in all phases of life. However, few participants believed that family predisposition [14, 15], unhealthy lifestyle, neglecting regular checkups [11, 13], and follow ups and destiny [15] were part of causes of their AMIs.

Some participants assumed that they were not at a risk of having AMI and disease would not affect them [13]. In one study participants believed that AMI only affected older population. In another study female participants did not consider themselves susceptible to AMI even those who had confirmed risk factors [16, 23].

Several participants believed that AMI would be preceded by warning sign and will give them time to change their life style [14]. Furthermore some participants were convinced by perception of their friends and families that AMI victims will have another second, third or even a fourth recurrent AMI. They assumed that their friends, neighbors who previously suffered from AMI were good sources of information [15]. Although few participants could identify some of the risk factors of AMI, additional education and information is much needed to educate the public regarding risk factors of AMI and the methods to reduce the modifiable risk factors.

Responses Related to the Event

According to the participants responses related to the event were affected by the various cause of AMI. They attributed their symptoms to ageing process, or to less serious causes like muscular pain, indigestion [20, 21, 25]. However, participants who previously had myocardial infarction sought medical attention sooner than other participants.²⁶ It

is clearly established that additional education will certainly help people increase awareness of AMI symptoms and timely response, so that they seek medical attention on time and as and when necessary [17].

Adopting Lifestyle Changes

Lifestyle change was most difficult challenge experienced by the patients. It was tougher than expected by the participants [14, 24, 25]. Some participants reported that fear, uncertainty and lack of knowledge made lifestyle changes more difficult. Few patients were worried that physical activity will do more harm and may lead to recurrent AMI attack [22]. On the other had some participants recognized the need to change their lifestyle in order to return to the activities they preferred to do before AMI attack [25, 26]. Some participants believed that lifestyle change were limited to their efforts to live within their capabilities.

Many participants considered surviving AMI as warning sign of impending death, which motivated them to make healthy lifestyle choices [14, 17]. For instance, some participants improved their nutritional habits, quit smoking and alcohol, meditation to decrease stress, working hour's reductions, engaging in recreational activities [15]. However, some participants performed strenuous activities too early which caused discomfort and discouraged them [19, 20]. Tailor made plans and individualized counseling may help survivors to develop appropriate lifestyle changes.

Family and Social Support

Majority of participants were satisfied with the professional support and treatment they received during hospitalization [24]. However, few of them stated that they have received better care and support after their discharge [25]. Most of the participants did not identify nurses as a source of information. In contrast, one group of participants identified nurses as major source of information [24].

The most stressful period for participants was the early period after being discharged. Patients needed more care and support during that time. Participants appreciated and valued the care and support of their family members in recovery process [14]. However, high stress among families related to their illness lead to hopelessness, frustrations in patients [11, 13]. Health care programs certainly need to provide additional support in early days and post discharge period. They also need to help and counsel family members/caregivers to learn appropriate support modalities.

Future Concerns and Insecurities

Many participants reported that they were keen to return to their previous lifestyle after surviving AMI [14], unfortunately, other participants were pessimistic about future; they were fearful of having another attack of AMI and its consequences [15]. They had a fear of death and saw life as fragile. Additionally participants also experienced feeling of helplessness, powerlessness and uncertainty regarding future [25, 26]. They believed that AMI has ended their productive and active life and some started preparing themselves for death. There was wide range of concerns for the future. Positive reinforcement, support and proper

information could help to allay fears and help them to tackle the challenges properly.

Expression of Needs

Participants expressed different needs which could be sorted in three phases: during hospitalization, discharge and follow up. Participants received general information like risk factors, healthy lifestyles during hospitalization. However they could not adapt during that period due to lack of knowledge about how to imply such changes. Patients were not prepared to hear the information they received, as they were overwhelmed emotionally. They also reported lack of directions from health professionals related to personalized plans and implementation of instructions [11, 12, 24].

All patients expressed the need for more professional support and information during discharge process. They needed information on how to deal with chest pain, manage stress, medications, nutritional guidance, healthy changes in lifestyles and cardiac rehabilitation programs.

Majority of participants felt need for better communication with health care providers during follow up period, after they settled back in their daily life [27, 28]. Since participants' had specific needs depending on their phase of recovery, health care professionals need to plan and provide time-sensitive educational teachings to patients when they are ready for it.

Discussion

The perspectives of people with heart disease about their knowledge on the nature and long term support they need not have been directly investigated till date; although it is recognized by all that they need long term support. To identify the desires and lived experiences expressed by participants is the main thrust of this analysis. This analysis includes aspects of long term monitoring and support the survivor needs as cardiac rehabilitation plan.

This qualitative literature review provided rich and fruitful data to explore AMI survivors' perspective and their personalized experiences. On one hand survivors were happy that they had survived, on the other hand they had to face a myriad of challenges as they worked to re-establish their lives, post AMI. Most of the participants lacked the ability to connect AMI with risk factors. Only few participants expressed dealing with modifiable risk factors to improve their quality of life. Other researchers also validated these concerns [29, 32]. Health care professionals need to plan community awareness campaigns to provide education about risk factors of the disease. Adoption of healthy lifestyle can modify risk factors like obesity, smoking, high blood pressure, stress, high cholesterol level, physical inactivity. The information related to healthy lifestyle can motivate people at risk to adopt lifestyle changes post myocardial infarction.

Participants who had experienced AMI were unable to recognize cardiac symptoms and attributed them to other causes. These findings were confirmed in study where AMI victims who delayed treatment and medical help attributed their symptoms to muscle pain, indigestion, flu like illness and breathing problems [33]. Studies have also found that women are less likely to recognize their susceptibility to AMI. These findings clearly indicate the need for continuous education for public, especially for women about the symptoms and to take proper steps while encountering these symptoms.

Survivors showed satisfaction in terms of role of family

throughout their recovery process, but they were not completely pleased with the support from health care professionals. Other researchers also have validated these findings [34, 35]. Health care providers should involve patients' families in educational campaigns and programs. This will orient and prepare families for managing survivors need appropriately and effectively. Discharge plan of patient should incorporate follow up plan, expectation of patients, professional resources as well. Moreover, additional support can be provided to survivors with the help of community health nurses by linking them with specialized community institutions. Nurses can also provide follow up information about the survivors. In India, nurses have the highest direct contact with the people. They could also be pioneers in health promotion and education programs.

Facing an AMI changed the participants' perspective towards life. Patients can be motivated to make appropriate lifestyle changes by using their positive feelings. Their pessimistic views can be remolded by using appropriate education and rehabilitation program to foster lifestyle changes.

Participants' needed appropriate information regarding hospitalization, post discharge and during follow up period. These phases are time sensitive and require different information and professional support to go through each phase comfortably. AMI survivors needed more professional support to resolve many issues like pain, fatigue, managing stress. They also required accurate information related to medications, nutritional guidance, and rehabilitation programs.

Limitations

Many journal articles are not available as full text, which limits access to potentially differing information.

Implications and Recommendations

Cardiac disease prevention and health promotion should be a priority for policy makers and clinicians as CVD is one of the leading causes of death globally. Additionally, health education related to cardiac diseases as well the proper management of responses should be included as public education campaigns. As the population of survivors from cardiac events increases, appropriate educational and rehabilitation program are required to be framed to meet their needs.

Nurses are the best resource person to help in making improvement in survivors' life. They can bridge the gap by providing information between survivors and information and planning educational campaigns. Furthermore, post discharge professional support should be provided to survivors with the utmost importance to the survivors needs.

Conclusion

It is important that patient's social environment should be considered as it can facilitate and hinder lifestyle changes in number of ways. This analysis has identified the need for personalized information regarding causes of illness, risk factors for illness and lifestyle changes. Finally, it is revealed that a one-size-fits-all approach is not applicable. Tailored based education has capacity to increase the perceived relevance of information accounts for better understanding. It is the need of the hour to have appropriate systems in place to facilitate the survival of AMI survivors. Surviving an AMI can be exacerbated by lack of

knowledge, lack of productive communication with health care staff, lack of follow up and professional support. Hence, there is a need of improvement in various domains like communication, modification of educational strategies and rehabilitation plans to ensure maximum benefits for the patients. Health care providers can use these reports to develop innovative pre and post AMI educational programs to improve survival and help the survivors to improve their post AMI lifestyles.

References

- Jayaraj CJ, Davatyan K, Subramanian SS, Priya J. Epidemiology of myocardial infarction, 2016(Accessed: August 17, 2019, at <http://dx.doi.org/10.5772/intechopen.74768>)
- Black C. Cardiovascular Diseases. WHO; [Last cited on 2016 Mar 13]. Available from: http://www.who.int/cardiovascular_diseases/en/
- Vahedian Azimi A, Alhani F, Ahmadi F, Kazemnejad A. The effect of family-centered empowerment model on the life style of the myocardial infarction patients. *IJCCN*. 2011; 2(4):127-32
- Gaziano TA, Gaziano JM. Epidemiology of cardiovascular disease. *Harrison's Principles of Internal Medicine*. 19th ed. New York, McGraw Hill; 216:266-65
- Gupta R, Gupta KD. Coronary heart diseases in low socioeconomic status subjects in India: an evolving epidemic. *Indian Heart J*, k gh, 2009; 61:358-67
- Global, regional and national comparative risk assessment of 79 behavioral, environmental and occupational and metabolic risks or cluster of risks, 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*, 2016; 388:1659-724.
- Gupta R. Burden of coronary heart disease in India. *Indian Heart J*, 2005; 57:632-8.
- Gupta R. Coronary heart disease in India: Absolute numbers and economic burden. Rapid response to Ghaffar A, Reddy KS, Singhi M. Burden of non-communicable diseases in South Asia. *BMJ*, 2004; 328:807. [PMC free article] [PubMed]
- Murray CJ, Lopez AD. Mortality by cause for eight regions of the world: Global burden of Disease Study. *Lancet*, 1997; 349:1269-76
- Polit D, Beck C. *Nursing research: Principles and Methods*. 7th ed, 2004. Lippincott Williams & Wilkins, 2004.
- Strauss A, Corbin J. *Basis of Qualitative Research: Grounded theory Procedures and Techniques*, 1990. Newbury Park. CA: Sage
- Kerr E, Fothergill-Bourbonnais F. The recovery mosaic: Older women's lived experiences after a myocardial infarction. *Heart Lung*. 2002; 31(5):355-367.
- Webster R, Thompson D, Mayou R. The experiences and needs of Gujarati Hindu patients and partners in the first month after a myocardial infarction. *Eur J Cardio Nurs*, 2002; 1:69-76.
- Bergan E, Bertero C. 'Grasp life again'. A qualitative study of the motive power in myocardial infarction patients. *Eur J Cardio Nurs*, 2003; 2:303-310.
- Jensen B, Petersson K. The illness experiences of patients after a first time myocardial infarction. *Pt Ed Couns*, 2003; 51:123-131.
- Clark A. 'It's like an explosion in your life: Lay perspectives on stress and myocardial infarction. *J Clin Nurs*, 2003; 12:544-553.
- Svedlund M, Danielson E. Myocardial infarction: narrations by afflicted women and their partners of lived experiences in daily life following an acute myocardial infarction. *J Clin Nurs*, 2004; 13:438-446.
- Hanssen T, Nordrehaug J, Hanestad B. A qualitative study of the information needs of acute myocardial patients and their preferences for follow-up contact after discharge. *Eur J Cardio Nurs*, 2005; 4:37-44.
- Gregory S, Bostock Y, Milburn BK. A qualitative study into lay experiences and the struggle to make lifestyle changes. *Family Practice*. 2006; 23(2):220-225
- Brink E, Karlson B, Hallberg L. Readjustment 5 months after a first-time myocardial infarction: reorienting the active self. *J Adv Nurs*. 2006; 53(4):403-411.
- Condon C, McCarthy G. Lifestyle changes following acute myocardial infarction: Patients perspectives. *Eur J Cardio Nurs*, 2006; 5:37-44.
- MacInnes J. The illness perceptions of women following symptoms of acute myocardial infarction: A self-regulatory approach. *Eur J Cardio Nurs*, 2006; 5:280-288.
- Johansson I, Swahn E, Stromberg A. Manageability, vulnerability and interaction: A qualitative analysis of acute myocardial infarction patients' conceptions of the event. *Eur J Cardio Nurs*, 2007; 6:184-191.
- Albarran J, Clarke B, Crawford J. 'It was not chest pain really, I can't explain it!' An exploratory study on the nature of symptoms experienced by women during their myocardial infarction. *J Clin Nurs*, 2007; 16:1292-1301.
- Kristofferzon M, LÖfmark R, Carlsson M. Striving for balance in daily life: experiences of Swedish women and men shortly after a myocardial infarction. *J Clin Nurs*, 2007; 16:391-401.
- Moonaghi KH, Mojalli M, Khosravan S. Psychosocial complications of coronary artery disease. *Iran Red Crescent Med J*. 2014; 16(6):e18162
- Christopher JM, Nicole DZ, Jane MH. A qualitative study of yomger men's experience of heart attack (myocardial infarction). 2017; 22(3):252-256
- Sharovsky LL, Romano BW, Franchini Ramires JA. Perception of depressive symptoms in patients after myocardial infarction: Qualitative study. *Heart Mind*, 2017; 1:107-11
- Nicolai J, Muller N, Noest S. *et al.* To change or not to change-That is the question: A qualitative study of lifestyle changes following acute myocardial infarction. *Chronic illness*. 2018; 14(1):25-41
- Ades PA. Cardiac rehabilitation and secondary prevention of coronary heart disease. *N Eng J Med*, 2001; 345:892-902.
- Dodani S, Mistry R, Farooqi M. *et al.* Prevalence and awareness of risk factors and behaviors of coronary heart disease in an urban population of Karachi, the largest city of Pakistan: a community survey. *J Pub Hlth*. 2004; 26(3):245-249.
- Jafary F, Aslam F, Mahmud H. *et al.* Cardiovascular health knowledge and behavior in patient attendants at four tertiary care hospitals in Pakistan-a cause for concern. *BioMed Central Pub Hlth*, 2005; 6:18.
- Khan M, Jafary F, Jafar T, Faruqi A, Rasool S,

- Hatcher J. *et al.* Knowledge of modifiable risk factors of heart disease among patients with acute myocardial infarction in Karachi, Pakistan: a cross sectional study. *BioMed Central Cardio Dis*, 2006; 6:18.
34. Eshah NF. Predictors of patients' delay to seek medical treatment among Jordanian acute myocardial infarction patients. JUST. (Unpublished Master Thesis), 2003.
 35. Simon T. Why is cardiovascular health important in menopausal women? *CLIMACTERIC*. 2006; 9(1 suppl.):s13-s18.
 36. Kristofferzon M, LÖfmark R, Carlsson M. Myocardial infarction: gender differences in coping and social support. *J Adv Nurs*. 2003; 44(4):360-374.