

International Journal of Medical and Health Research



Volume: 1, Issue: 2, 23-25
Sep 2015
www.medicalsjournals.com
ISSN: 2454-9142

Namama Soran Talabani
Chemistry department,
Faculty of Science,
University of Sulaimania.

Monitoring Levels of Iron, TIBC, Hb, Transferrin and Ferritin during Pregnancy Trimesters and Lactation in Sulaimania City /Iraq

Namama Soran Talabani

Abstract

Iron deficiency is the most common nutritional disorder in the pregnant women. This study was conducted to investigate the possible contribution and impact of iron status on pregnancy in a Sulaimnia/Iraq population. Two hundred and thirty five pregnant women aged 18-43 years. The women were followed up weekly till delivery after which neonatal anthropometrics and other birth out comes were recorded. Mean serum Hb, ferritin and iron were found to be lower in 3rd trimester as compared to 1st trimester, which become lower at postpartum then reach nearly to normal value during lactation. While serum ferritin was found to be lower in 2nd trimester as compared to 1st trimester, it was significantly higher in 3rd as compared to second trimesters, it is also decreased at postpartum and lactation. Serum TIBC, transferrin and Fe/TIBC levels increased during 1st, 2nd and 3rd trimesters, Serum TIBC also elevated at postpartum, then decreased during lactation. This result provides that, it is important to detect which form of iron become abnormal during trimesters of pregnancy, postpartum and lactation which may lead to cause of nutritional deficiency in pregnant women, so as to reduce public problem.

Keywords: Pregnancy, iron form, postpartum and lactation.

1. Introduction

Iron is one of the most important micronutrients for human populations, given its central role in key biological processes. Iron is especially critical during pregnancy given the rapid cell and tissue development involved in fetal growth. Pregnancy has a net iron cost in the range of 600-800mg^[1, 2].

It has been reported that 56% OF pregnant women in low income countries are affected in contrast to 18% in high income countries^[3]. In pregnant women, an adequate iron status is important to ensure an uncomplicated pregnancy as well as a normal development of the fetus and maturity of the newborn child^[4].

Iron deficiency during pregnancy is associated with a number of maternal and fetal problems including the risks of preterm births, low birth weight babies, perinatal mortality and intrauterine growth retardation^[5]. The hemoglobin concentration is often used as a pseudo marker for iron deficiency. However, hemoglobin is not suitable to asses iron-status especially not in pregnancy due to hypervolaemia and haemodilution. Hemoglobin yields information about the presence of anemia in general when body iron reserves are depleted^[6].

The diagnostic workup for iron deficiency includes red blood cell indices, serum iron, serum total binding capacity, serum transferrin saturation and serum ferritin level. Serum iron transferrin is frequently abnormal in pregnancy. Low sensitivity of transferrin saturation and day to day and even hour to hour fluctuation of serum iron levels renders it less efficient than serum ferritin level for diagnosing iron deficiency which is the only condition associated with decreased serum ferritin concentration^[7, 8]. Serum ferritin, a marker of iron storage, has the advantage of being a sensitive indicator of iron deficiency, but because it is increased in the presence of inflammation, ferritin is not a specific indicator of iron deficiency^[9, 10, 11].

Approximately 800 mg of iron are requirement in pregnancy, over and above the 230 mg of iron that the women would have required even if she had not been pregnant, and the 150mg that she may lose through blood loss at delivery^[12, 13]. Though this iron expenditure is offset by the lack of menstrual blood loss during this time, the net cost is still high enough that iron recommendation during pregnancy (27 mg/day) for exceed those for non-pregnant, Not

Correspondence
Namama Soran Talabani
Chemistry department, Faculty
of Science, University of
Sulaimania.

Lactating women (18mg/day) (1). While iron recommendations for lactating women are much lower than those for non-pregnant, non-lactating women (9mg vs. 18mg), this number is based on the assumption of lactation induced amenorrhea and does not take into account that many women enter or conclude pregnancy with iron insufficiency or deficiency [1].

The aim of the present study therefore is to report the values of some important iron forms as parameters in apparently healthy pregnant Iraq subjects residing in kurdistan region (sulaimania City). The parameters assessed include: serum iron (Fe), total iron binding capacity (TIBC), serum transferrin, serum ferritin, hemoglobin and Fe/TIBC%. These parameters was study also during postpartum and lactation which compared with normal women. This hope would further our attempts at establishing normal reference ranges for these parameters for pregnant women in Sulaimania/Iraq.

2. Materials and Methods

This prospective, comparative cross-sectional study was performed at Sulaimania maternity hospital with collaboration of college of science, chemistry department, Sulaimania University from October 2014 to march 2015.

A total number of 235 apparently health pregnant subjects aged between 18-43 years were recruited into the study. All subjects were selected randomly during their antenatal visits. Patients with history of Diabetes, hypertention, hepatitis and any other acute or chronic illness were excluded from the study.

A 2-3 ml of EDTA blood sample was collected for hematological parameters which were measured on a fully

automated hematology analyzer (systmex KX-21) and 3 ml of blood was collected in gel tubes. Blood was centrifuged at 3000 rpm for 5min. the supernatant was used to quantify the serum TIBC and iron level by spectrophotometric methods supplied by Biolabo, France. The ceruloplasmin levels in human serum by turbidimetric method supplied fortress Diagnostics [14].

Transferrin can be estimated indirectly from the TIBC value by the following equation [15].

$$\text{Transferrin (g/dl)} = 0.7 \times \text{TIBC (g/dl)}$$

All statistical analyses in studies were performed using SPSS version 17.0 for

Windows (Statistical Package for Social Science, Inc., Chicago, IL, USA). Descriptive analysis was used to show the mean and standard deviation of variables.

3. Results

A total (280) women were selected in the study, (235) pregnant women as test group and (45) non-pregnant women as control. overall, the subjects were in the age range 18 – 43 years. Age distribution of the studied group was between 18 to 43 years old, and that of the pregnant group irrespective of the trimester of pregnancy and the lactation was comparable to that of the control group $p > 0.05$.

Table (1) shows the age representation, number of abortion and number of children in pregnant women in three trimesters and at postpartum with lactation.

Table (2) shows: serum iron, serum ferritin, total iron binding capacity, hemoglobin, serum transferrin and Fe/TIBC%.

Table 1: Revelant Clinical Parameters

Parameters	Control No of case	1 st trimester	2 nd trimester	3 rd trimester	P.P	lactation
Age Years	19-42	18-38	18-39	19-41	18-40	19-41
No of abortion	0-1	0-1	0-3	0-1	0-2	0
No of children	0-2	1-3	0-3	0-1	1-2	0-2

Table 2: Comparison of pregnant women classified by trimester of pregnancy and lactation with control subjects

Parameters	control	1 st trimester	2 nd trimester	3 rd trimester	Postpartum	lactation
S. Iron g/dl	70.15±12.92	60.93±14.23	51.97± 14.20	50.99± 18.89	48.89± 16.78	73.25± 10.83
Ferritin ng/ml	53.95± 15.12	17.02± 4.8	14.69± 5.9	16.01± 4.91	14.09± 3.37	13.01± 2.91
TIBC gm/dl	300.86± 25.13	320± 95	339.81± 49.93	442.06± 41.04	472.5± 80.06	390.91± 21.82
Hemoglobin gm/dl	12.43± 2.18	11.81 ±0.87	10.91± 1.01	10.38 ±0.69	9.92± 0.76	9.80±0.23
Transferrin mg/dl	200.1± 11.9	204.81± 12.82	239.81± 19.37	225.19± 20.09	290.31± 19.83	295.81 ±20.31
Fe/TIBC%	23.3	18.9	15.3	11.53	16.84	18.73

Discussion

Pregnancy is a risk factor for development of iron deficiency anaemia. In developing countries iron deficiency is very common, even without pregnancy. This study was designed to assess iron status in Sulaimanian pregnant women at the time of delivery, regardless of whether they were taking iron supplementation or not, in order to find out the prevalence of iron deficiency among them, we also take serum from women during lactation in order to find which one of these markers affected by lactation.

Statistically significant difference were observed in the values of serum iron concentration amongst the three trimesters of pregnancy, highest value being observed amongst subjects in the first trimester and lowest value amongst subjects in the third trimesters of pregnancy. This is consistent with reports of changes in serum iron concentration during pregnancy in

Filipino, Caucasin and Tariah [16-18].

During pregnancy the physiologic need for iron is extraordinary high. In women of reproductive age, the requirement for absorbed iron is on average 0.8mg/day. The average requirements for absorbed iron increased steadily during pregnancy from 0.8mg/day in the third trimester. The average requirement for absorbed iron in the entire gestation period is ~4.4 mg/day [19, 20].

This study agreed with Fenton *et al.* in 1997 who had concluded that serum iron and TIBC are frequently abnormal in pregnancy and so they are of little help in diagnosis of iron deficiency anemia during pregnancy [21]. Serum ferritin was found to be low in most mothers ferritin level is the best parameter for assessment of iron are unreliable during pregnancy [22]. During pregnancy Hb levels in woman is naturally lower than when she is not pregnant. This is because

the fluid (plasma) increases by about 50% during pregnancy (peaking at about 32 weeks) [23]. The increased plasma dilutes the red cells, making their level drop.

Results of this study did not differ from those of other studies carried out in different parts of the country and abroad and have been found in conformity with previous works carried out by various workers. Several factors were observed to be responsible for a high rate of iron forms deficiency in a community like ours. Multi parity, poor socio economic and educational status are the principal reasons for a high prevalence of iron deficiency anemia in our population [24].

The aim of this paper is to describe the consequence of iron forms deficiency in pregnancy and to outline strategies for prevention.

Reference

- Picciano MF. Pregnancy and lactation: physiological adjustments, nutritional requirements and the role of dietary supplements, *J Nutr.* 2003; 133:1997S-2002S.
- Gambling L, Lang C, Mc Adrdle HJ. Fetal regulation of iron transport during pregnancy, *Am J Clin Nutr.* 2011; 94:190S-1907S.
- United Nation Administrative committee on coordination, Sub-committee on Nutrition (ACC/SCN). Fourth report on the World Nutrition Situation: Nutrition throughout the life cycle. Geneva: ACC/SCN in collaboration with international food policy research Institute, 2000.
- Milman N. prepartum anemia: prevention and treatment. *Ann Hematol* 2008; 87:949-951.
- Saeed GA, Khattak N, Hammid R. Anemia in pregnancy and spontaneous preterm birth, *J Pakistan Inst Med Sci* 2002; 13(2):698-701.
- Milman N. Iron and pregnancy a delicate balance: *Ann Hematol* 2006; 85:559-565.
- Naghmi A, Khalid H, Shaheen M. comparison of serum ferritin levels in the trimesters of pregnancy and their correlation with increasing gravidity, *Inter J of pathology.* 2007; 5(1):26-30.
- Fal TK, Lao TT. Hemoglobin and red cells indices correlated with serum ferritin concentration in late pregnancy. *Obstetrics and Gynecology* 1999; 93:427-431.
- WHO Assessing the iron status of populations. Including literature reviews. Report of a Joint World Health Organization /Centers for Disease Control and prevention technical consultation on the Assessment of Iron Status at the population level. World Organization, Geneva, Switzerland, 2004.
- Zimmermann MB. Methods to assess iron and iodine status, *Br J Nutr.* 2008; 99(3):S2-S9.
- Iron status Indicator. Available online: http://www.cdc.gov/nutrition_report/99-02/pdf/nr-ch3.pdf (accessed on 24 may 2014).
- Bothwell TH. Iron requirements in pregnancy and strategies to meet them. *Am J clin Nutr.* 2000; 72:265S-271S.
- Bender DA. Do we really know vitamin and mineral requirement, for infants and children? *J R Soc Promot health.* 2003; 123:154-158.
- Tietz NW. In Text book of clinical chemistry, edited by N W Tietz WB. Saunders Company, Philadelphia, London, Toronto, 960-962.
- Dacie JV, Lewis SM. Practical Hematology. 7th edition Edinburgh Churchill Livingstone.
- Africa KMC, Aghana LE, Alday A. A preliminary determination of serum transferrin levels in normal Filipino pregnant women in the three trimesters of pregnancy, *Acta Medica philippina* 1997; 202(90):120-126.
- Kaneshige E. Serum ferritin as assessment of iron stores and other hematologic parameters during pregnancy. *Obstet Gynaecol* 1981; 57(2):238-242.
- Amah Tariah FS, Ojeka SO, Dapper DV. Hematological values in pregnant women in Port Harcourt, Nigeria II: Serum iron and transferrin, total and unsaturated iron binding capacity and some red cell and platelet indices. *Niger J physiol Sci.* 2011; (26):173-178.
- Bothwell TH. Iron requirements in pregnancy and strategies to meet them. *Am J Clin Nutr.* 2000; 72:257S-264S.
- Milman N. Iron and Pregnancy delicate balance. *Ann Hematol* 2006; 85:559-565.
- Harthoorn Lasthuizen EJ, Lindemas J, Lindemas J, Langenhuijsen MM. Does iron deficient erythropoiesis in pregnancy influence fetal iron supply? PIMD 40968569 *Acta Obst Gynecol Scand* 2001; 80(5):392-6.
- Enaam A, Abdegader, Tayseer AD, Anwar A, Sana EA, Haemoglobin level, RBCs indices, and iron status in pregnant females in Sudan, *Basic Research Journals.* 2014; 3(2):08-13.
- Baynes RD, SKikne B. Iron absorption. In Brock, JH. Halliday JW. piparred MJ, powell L W(eds). *Iron metabolism in health and disease.* London, UK WB Saunders 1994; 151-87.
- Ijlal F, Jalali S, Khan NB, Khan AQ. Prevalence of anemia in women at District head quarter Hospital Gilgit, *Pak J Med Res.* 2000; 39:70-80.