



Clinical Profile and laboratory indicators in children on admission based Predicting the severity of dengue fever: A study in Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh

Dr. Mohammad Mashiur Rahman¹, Dr. Md. Shafiu Hoque², Dr. Asma Hoque³, Dr. Mohammed Akter Hossan Masud⁴

¹ Registrar, MBBS, DCH, MPH, Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh

² MD. Associate Professor of Pediatrics, Bangladesh Institute of Child Health, Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh

³ Resident Medical Officer, MBBS, DCH, MPH, Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh

⁴ Resident Medical Officer, MBBS, Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh

Abstract

Introduction: Dengue fever is a re-emerging viral disease commonly occurring in tropical and subtropical areas. The clinical features and abnormal laboratory test results of dengue infection are similar to those of other febrile illnesses; hence, its accurate and timely diagnosis for providing appropriate treatment is difficult. Delayed diagnosis may be associated with inappropriate treatment and higher risk of death. Early and correct diagnosis can help improve case management and optimise the use of resources such as hospital staff, beds, and intensive care equipment. The goal of this study was to develop a predictive model to characterise dengue severity based on early clinical and laboratory indicators using data mining and statistical tools.

Objective: To find out clinical Profile and laboratory indicators in children on admission based Predicting the severity of dengue fever.

Methods: We retrieved data from a study of febrile illness in children in Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh Of 325 febrile episodes recorded, 198 patients were confirmed to have dengue. A classification and regression tree (CART) was used to construct a predictive decision tree for severe dengue, while logistic regression analysis was used to independently quantify the significance of each parameter in the decision tree.

Results: This study was 325 patient admissions during the study year, of which 251 (77.23%) met the inclusion criteria. Clinical manifestations of 198 patients with dengue infection, including 38 with severe disease. Of these, 74 (22.76%) were not enrolled, leaving 125 febrile episodes in 180 children, with 144 children having a single episode, 31 children having two episodes, one child having three episodes, and four children having four episodes. A decision tree algorithm using haematocrit, Glasgow Coma Score, urine protein, creatinine, and platelet count predicted severe dengue with a sensitivity, specificity, and accuracy of 60.5%, 65% and 64.1%, respectively.

Conclusions: The decision tree we describe, using five simple clinical and laboratory indicators, can be used to predict severe cases of dengue among paediatric patients on admission. This algorithm is potentially useful for guiding a patient-monitoring plan and outpatient management of fever in resource-poor settings.

Keywords: classification tree, dengue, severity, Cambodia, data mining, children

1. Introduction

Dengue fever causes a high burden of disease and mortality across tropical and subtropical regions in Southeast Asia, Africa, the Western Pacific, and the Americas [1]. Dengue virus comprises four serotypes, DENV-1, DENV-2, DENV-3, DENV-4 which are transmitted by *Aedes aegypti* mosquitoes [2-4]. An estimated 2.5 billion people worldwide are at risk of dengue. More than 50 million dengue infections are estimated to occur annually, of which approximately 500,000 result in hospital admissions for severe dengue in the form of dengue haemorrhagic fever (DHF) or dengue shock syndrome (DSS), principally among children [5]. Dengue infection is frequently confounded with other febrile illnesses (OFI), presenting with non-specific clinical symptoms and clinical features analogous to OFI. During the early stages of dengue, the presence of non-specific febrile illness makes precise diagnosis strikingly difficult, resulting in inefficient treatment and possible increases in morbidity and mortality [2, 6]. Severe dengue fever, if not appropriately managed, may lead to rapid death,

particularly in children [7, 8]. In addition, the lack of necessary laboratory facilities, particularly in remote, rural areas, may cause difficulty in discriminating dengue infection from OFI [9]. Dengue is one of the most common vector-borne diseases in Southeast Asia, and one of the most important mosquito-borne viral diseases with an epidemic potential in the world [10]. Dengue was first included in Cambodia's national surveillance programme in 1980. Since 2000, between 10,000 and 40,000 dengue cases have been reported annually by the Dengue National Control Program [11], from a total population of approximately 13.5 million people [12]. The true incidence of the burden of disease in Cambodia remains under-reported due to difficulties in diagnosing dengue infection, especially in hospitals [13]. In this study, data from a cohort of children that were admitted with febrile illness to Angkor Hospital for Children, Siem Reap, Cambodia, during a one-year period were retrospectively analysed using a data mining approach. This approach used a classification and regression tree, or CART, which was first introduced by Breiman *et al.* [14]

This is a common tool used in data mining, which creates a model or algorithm that predicts the value of a target variable based on several input variables. In our study, CART was constructed to predict the severity of dengue infection based on early clinical and laboratory indicators. The model was then evaluated against the final diagnoses.

2. Objective

- a) To find out clinical Profile and laboratory indicators in children on admission based Predicting the severity of dengue fever.

3. Methods

We conducted a retrospective study of data derived from an investigation of febrile illness in children (“the fever study”) in Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh. We retrieved data from a study of febrile illness in children in Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh Of 325 febrile episodes recorded, 198 patients were confirmed to have dengue. A classification and regression tree (CART) was used to construct a predictive decision tree for severe dengue, while logistic regression analysis was used to independently quantify the significance of each parameter in the decision tree. This is in Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh, which provides free, comprehensive healthcare to children aged less than 16 years of age, and includes specialised medical and surgical in-patient and outpatient care. For the fever study, the inclusion criteria were age < 16 years, documented axillary temperature ≥ 38.0 °C within 48 h of admission, and informed consent by a parent or caregiver. Children who developed fever ≥ 48 h after admission or following surgery were excluded since they could be considered as having acquired a healthcare-associated infection [16, 17]. Integrated Management of Childhood Illness (IMCI) was used for the assessment and decision-making on whether to admit a patient to the hospital [18]. Data were collected on admission by clinicians using a specific case report form. Admission blood samples and, where possible, a convalescent serology sample taken on discharge, or seven days after admission, were taken for IgM antibody and NS1 antigen testing. All admitted patients were reviewed twice daily for eligibility and data collection quality. Data were collected between 12th October 2017 and 12th October 2018 from patients who were admitted. Patients were classified as having dengue virus infection if NS1 antigen was detected in their serum by ELISA, or if paired sera from acute and convalescent time points (≥ 7 days following the acute sample) showed rising or static anti-dengue IgM (and anti-dengue IgM was greater than anti-Japanese encephalitis IgM) [15]. NS1 antigen and IgM antibody results were combined in a Boolean manner using AND/OR operators to ensure that the entire temporal spectrum of patient presentations during the acute phase of dengue infection were covered, with NS1 antigen present in the serum in the early phase of infection and dengue IgM antibodies usually present after 2–5 days of infection [19]. The ratio of anti-dengue to anti-JEV IgM levels was used to determine if the infection was dengue or Japanese encephalitis virus, whose antibodies often have some cross-reactivity when co-circulating in the same area. Children less than 60 days old were not tested for dengue virus infection. All confirmed dengue cases were further categorised as either severe or non-severe dengue. From our literature review we noted

that, although the revised 2009 WHO classification was said to be an improvement on the 1997 WHO classification, there was still a need for training, dissemination of relevant information, and further research on the warning signs of severe dengue [20]. The classification was also considered by many to be too broad, requiring more specific definitions of the warning signs [21], that it increased the workload for health care personnel, and was not simple or user-friendly enough [22]. In our study we categorised dengue cases as severe based on a two-step process. The first step was to take into account all confirmed dengue cases with intensive care unit (ICU) admission, together with the 2009 WHO dengue classification. Secondly, two independent pediatricians’ assessments were considered to a) exclude any ICU-admitted cases that might not have had severe dengue as their primary diagnosis and b) to include any non-ICU-admitted cases which may have actually presented with severe dengue but were not admitted for some reason, usually because of resource limitations. The presence of bleeding was not assessed other than by examining stool samples for blood. The case-by-case assessment and verification by the two clinicians was used as the reference for the predictive model.

Data analysis and construction of a predictive model

The demographics and clinical characteristics of severe and non-severe dengue cases were described using the mean \pm standard deviation (SD) if the data were normally distributed, or by median and range otherwise. Comparisons between the two groups were performed using the Student’s t-test for continuous variables if the data were normally distributed, otherwise the Mann–Whitney U test was used. A chi-square test was used for categorical data. A p value < 0.05 was considered significant. A classification and regression tree (CART) was constructed for predicting the severity of dengue cases based on their early clinical features and laboratory indicators on admission. The J48 algorithm was used for generating decision trees because it is able to handle nominal, categorical and numerical data, as well as missing values. The loss matrix was specified to differently weight misdiagnoses. In this study we assigned five times greater weight to false negatives when compared with false positives, i.e. the cost of misdiagnosing a patient with severe dengue was five times greater than a non-severe case being misdiagnosed as severe. Pruning and tuning parameters were applied to optimise the predictive model by avoiding an over-complex tree, and thus increasing the model’s accuracy. The 10-fold cross-validation function provided by Weka was used to estimate the out-of-sample accuracy, given the constraint on data availability and avoiding the over-fitting issue. Put simply, it split the data set into ten partitions, nine of which were for training, with one partition for testing. The tree model was built on the training set, and applied to the testing set. To reduce variability, multiple rounds of cross-validation were performed using different partitions, and the validation results were combined over the rounds to estimate the model’s performance [23]. Once the final tree was obtained, the significance of each predictive factor was then quantified through multiple logistic regression with the ‘enter’ method of selection (i.e. all variables were included in the model) and reported as an odds ratio (OR) and 95% confidence interval (95% CI). Descriptive analysis and multiple logistic regression were performed using the

Statistical Package for the Social Sciences (SPSS) software, version 13.0.

Parameterisation

Before data mining algorithms can be used, a target data set must be assembled and pre-processed, which involves cleaning, removing, grouping, and transforming the data. There were 24 variables originally available for the analysis. However, three variables were excluded from the analysis i.e. tourniquet test result was with more than 15% missing data points where as pulse rate and respiratory rate were age-dependent parameters. The latter ones were excluded from the analysis since they would not be practical to refer to if presented in the final model. For other variables with fewer than 15% missing values, the missing values were imputed using a one-by-one single imputation approach. The advantage of the imputation method over the tree-based mining algorithm within CART [24] is that it separates the missing data problem from the prediction problem, allowing different predictive modelling methods to be applied to the imputed data set [25]. In our study, some missing values were imputed with a single value, including the mean value for some variables (number of days of fever, capillary refill time, Glasgow Coma Score, and urea result) and the median value for others (haematocrit, creatinine, ALT, respiratory rate of infant, urinary protein and RBC, and WBC, neutrophil, lymphocyte, and platelet counts).

4. Results

This study was 325 patient admissions during the study year, of which 251 (77.23%) met the inclusion criteria. Clinical manifestations of 198 patients with dengue infection, including 38 with severe disease. Of these, 74 (22.76%) were not enrolled, leaving 125 febrile episodes in 180 children, with 144 children having a single episode, 31 children having two episodes, one child having three episodes, and four children having four episodes. The patients were mainly diagnosed as having lower respiratory tract infection (38.3%), undifferentiated fever (25.5%), or diarrhoeal disease (19.5%) [15]. Out of 180 enrolled children, there were 19 deaths, the causes of which were: clinical pneumonia with no organism/virus identified (12 cases, 27.5%), dengue virus infection (eleven cases, including one with co-existent melioidosis, two with co-existent scrub typhus, and four with coexistent clinical pneumonia, 15.9%), and melioidosis (four cases, 5.8%). 41 non-dengue episodes and 86 episodes with no samples available were excluded from this analysis. Further details can be found in the original report [15]. Out of 198 confirmed dengue episodes, 43 episodes required ICU admission, with 29 of those classified as severe dengue based on their clinical signs, supported by two independent clinical opinions. Nine additional severe dengue episodes were included from non-ICU admissions, making a total of 38 episodes of severe dengue. There were eleven in-hospital deaths amongst all ICU-admitted patients with dengue virus infection, however dengue was the primary diagnosis in just five of these. Therefore, only these five cases were included in the severe dengue group. The flowchart of the study is shown in Fig. 1. The three most common features among patients with severe disease were ICU admission (76.3%), rapid respiratory rate (81.5%), and rapid pulse (65.7%). Severe

dengue was more prevalent in children aged less than five years old. Vomiting and abdominal pain were significantly more common in the severe dengue group, as were rapid pulse and respiratory rate, increased capillary refill time, and low GCS. A significantly higher proportion of patients with severe dengue presented with a lower hematocrit, higher WBC and lymphocyte count, higher ALT level, together with the presence of urinary RBC (Table 2). The final decision tree algorithm included five clinical and laboratory parameters: hematocrit, GCS, urinary protein, creatinine, and platelet count. The sensitivity and specificity of the model were 60.5% and 65%, respectively (Fig. 2). The accuracy of the model was 64.1% where the clinical diagnosis was used as the reference value. The area under the receiver operating characteristic (ROC) curve for logistic regression was 0.616. The final decision tree was then restructured using logistic regression analysis to estimate the impact of each CART-selected variable as represented by the OR and 95% CI. Table 3 gives the estimated OR for each parameter selected by CART. Low hematocrit, low GCS, low platelet count, presence of urine protein, and high creatinine increased the probability of a diagnosis of severe dengue, with significant OR ranging from 1.47 to 13.73. This study did not specifically focus on children and included only female patients. The sensitivity and specificity achieved in its decision algorithm were 81% and 54%, respectively.

Table 1: Clinical features and laboratory indicators of 38 severe dengue cases based on the 2009 WHO dengue classification.

Factor	N (%)
ICU admission	29 (76.3)
Average number of ICU days	4.6 days
Vomiting	15 (39.4)
Blood in stool	(21)
Rapid respiratory rate	(81.5)
Rapid pulse	25 (65.7)
Liver enlargement	22 (57.8)
Abdominal pain	(23.6)
Low WBC count	10 (26.3)
High WBC count	13 (34.2)
Low haematocrit	22 (57.8)
High haematocrit	(7.8)
Low platelet count	12 (31.5)
Low Glasgow Coma Scale	(21)
High urea	(23.6)
High creatinine	21 (55.2)
ALT > 1000	(5.2)
Urine protein > 100	(13.1)
Urine red blood cells	12 (31.5)
Death	(13)

Clinical features, including blood in the stool, liver enlargement, ICU admission, number of days in ICU, low or high hematocrit, low or high WBC count, high creatinine, high urea, low platelet count, rapid pulse, rapid respiratory rate, low Glasgow Coma Score (GCS), pleural effusion (only one case), abdominal pain, urinary protein, urinary RBC, and high ALT, were considered on a case-by-case basis when clinicians classified dengue as severe or non-severe. The clinical features and laboratory indicators of the 38 severe dengue cases are shown in Table 1.




Fig 1: The flowchart of the study. Boxes show the total number of patients enrolled in the study, reasons for exclusion from the analysis, model construction and evaluation.

There were eleven in-hospital deaths amongst all ICU-admitted patients with dengue virus infection, however dengue was the primary diagnosis in just five of these.

Therefore, only these five cases were included in the severe dengue group. The flowchart of the study is shown in Figure 1.

Table 2: Clinical manifestations of 198 patients with dengue infection, including 38 with severe disease.

Variables	Non-severe (n = 160)	Severe (n = 38)	p value
Demographics			
Male (n, %)	84 (52.5)	23 (60.5)	0.372
Age: 28 days to 1 year (n, %)	40 (25.0)	16 (42.1)	
≥ 1 year to < 5 years	45 (28.1)	14 (36.8)	0.012
≥ 5 years to < 16 years	75 (46.9)	8 (21.1)	
History/symptoms			
Number of days of fever	4.31	4.10	0.683
Vomiting (n, %)	94 (58.7)	15 (39.4)	0.032
Abdominal pain (n, %)	75 (46.8)	9 (23.6)	0.012
Headache or retro-orbital pain (n, %)	63 (39.3)	8 (21)	0.101
Clinical parameters			
Rash (n, %)	18 (11.2)	1 (2.6)	0.153
Temperature (°C)	38.76	38.63	0.234
Pulse/min	131.49	152.26	< 0.001
Capillary refill time	2.03	2.24	0.001
Respiratory rate	38.40	46.42	0.002
Liver enlargement (n, %)	64 (40.0)	22 (57.8)	0.127
Glasgow Coma Score	14.60	13.42	0.003
Laboratory parameters			
Haematocrit (%)	32.56	28.84	0.004
Platelets (per 10 ³ /µl)	267.87	294.16	0.477
White blood cells (per10 ³ /µl)	9.46	14.25	0.006
Neutrophils	5.64	8.01	0.46
Lymphocytes	2.76	5.14	< 0.001
Urea (mmol/L)	4.64	4.95	0.712
Creatinine (µmol/l)	68.29	68.71	0.932
Alanine transaminase, IU/l	78.99	177.08	0.002
Urine protein mg/dL	12.34	24.13	0.088
Urine red blood cells	2.84	10.32	0.009




Fig 2: Clinical decision tree to distinguish severe dengue from all cases of dengue (HCT = haematocrit, GCS = Glasgow Coma Score, PLT = platelets).

Using a data mining approach, we have developed an algorithm using both simple clinical manifestations and laboratory indicators to predict the severity of dengue during the early phase of the illness. The final algorithm for predicting severe dengue (Figure 2) comprised six components in order of their significance.

Table 3: Output from logistic regression using the decision tree algorithm to predict severe dengue infection.

Variable	OR	95% CI		p value
		Lower	Upper	
Haematocrit (> 28%)	1			
Haematocrit (≤ 28%)	7.114	3.000	16.869	< 0.001
Glasgow Coma Score (> 11)	1			
Glasgow Coma Score (≤ 11)	13.731	3.460	54.501	< 0.001
No urine protein	1			
Urine protein	1.832	0.777	4.319	0.167
Creatinine (≤ 84 µmol/l)	1			
Creatinine (> 84 µmol/l)	1.471	0.509	4.247	0.476
Platelets (> 146 × 10 ³ /µl)	1			
Platelets (≤ 146 × 10 ³ /µl)	2.334	0.946	5.763	0.066

The parameters that were statistically associated with severe dengue were 1) low haematocrit (OR = 7.114, 95% CI = 3.00–16.87, p < 0.001) and 2) low GCS (OR = 13.73, 95% CI = 3.46–54.50, p < 0.001). Although low platelet count (OR = 2.33, 95% CI = 0.95–5.76), presence of urine protein (OR = 1.83, 95% CI = 0.78–4.32) and increased serum creatinine (OR = 1.47, 95% CI = 0.51–4.25) were associated with an increased risk of severity, they were not shown to be statistically significant by regression analysis (Table 3).

5. Discussion

Using a data mining approach, we have developed an algorithm using both simple clinical manifestations and laboratory indicators to predict the severity of dengue during the early phase of the illness. The final algorithm for predicting severe dengue (Figure 2) comprised six components in order of their significance. The most significant factor in predicting severe dengue was low haematocrit, followed by a GCS of 11 or below as the second split if haematocrit was greater than 28, the presence of urine protein and creatinine above 84 µmol/l as the third split if GCS was above 11, and finally a platelet count of 146,000 per mm³ or less as the final split, if the presence of urine protein and creatinine was below 84 µmol/l.

Comparing the algorithm we derived with those re-reported in previous studies, we found both similarities and differences. Potts *et al.* constructed decision algorithms for predicting dengue shock syndrome (DSS) or dengue with significant pleural effusion [26]. The algorithm achieved a high sensitivity of 97%. Both low haematocrit and platelet counts were also identified as predictive factors in their work, although the cut-off values used in our algorithm were more extreme, i.e. for haematocrit ≤ 28 vs. ≤ 40, and for platelet count ≤ 146,000 vs. ≤ 160,200. The mechanism by which thrombocytopaenia is caused by dengue virus is complex [27]. Previous studies suggest that the virus probably contributes to bone marrow suppression and platelet destruction [28, 29]. To meet the WHO guidelines for classifying patients with DHF, thrombocytopaenia (platelet count ≤ 100,000) is re-quired. Srikiatkachorn *et al.* demonstrated that thrombocytopaenia was related to dengue severity and that not all severe cases would have been classified as DHF according to the WHO criteria [30]. Although thrombocytopaenia suggests that dengue infection is severe, a low platelet count is also common among OFI such as malaria and scrub typhus [31]. The 1997 WHO definition of DHF stated that a low platelet count (≤ 100,000), together with an increased haematocrit of ≥ 20% above the baseline value, is indicative of plasma leakage. In contrast, our re-sults and those of Potts *et al.* suggested a drop in haem-atocrit as a sign of severity, especially among patients with internal bleeding in areas such as the gastrointestinal tract [26]. Our results also suggested a more extreme haemato-crit value compared with the previous study (28% vs 40%) [27]. Although Potts *et al.* identified WBC count and monocyte percentage as important, our analysis did not identify monocyte results as significant even when in-cluded in the decision tree algorithm. In addition, Potts *et al.* evaluated predictors for DSS and dengue with significant pleural effusion, whereas in our study severe dengue was differentiated based on clinical features and laboratory indicators. Another recent study, by Tamibmaniam *et al.*, used simple logistic regression and identified three parame-ters, including vomiting, pleural effusion, and low sys-tolic blood pressure, to predict severe dengue based on the 2009 WHO criteria [32]. This study did not specific-ally focus on children and included only female patients. The sensitivity and specificity achieved in its decision algorithm were 81% and 54%, respectively. Of the three parameters they identified, vomiting was the only parameter available in our study, and although it initially appeared to be significant in the severe group, it was not selected for the final tree. Despite using a similar approach to predict somewhat similar outcomes to the aforementioned studies, we identified additional parameters that related to the sever-ity of dengue, including GCS, urine protein, and serum creatinine. There are a number of possible explanations for these differences, as outlined below. GCS is used to measure the level of consciousness (mental status changes) [33]. In our results, the node with GCS ≤ 11 (considered to be moderate) in the model was significant. Rao *et al.* showed that patients with den-gue encephalitis had a GCS of 7–8 and recommended intubation and mechanical ventilator support during their hospitalization [34]. Previous studies in which urine protein was associated with DHF or DSS used the urine protein-to-creatinine ratio [35, 36], but we used only a urine dipstick for this measure. The presence of urine protein in severe dengue could be due

to plasma leakage. An increased serum creatinine level indicates kidney dysfunction. In patients with DHF, a mild increase in serum creatinine is common, in contrast to the higher levels seen in severe dengue cases. Our model showed that a serum creatinine level > 84 mmol/l (4.6 mg/dl) was associated with severe dengue, a value similar to that found in Thai paediatric patients with DHF, whose mean serum creatinine was 4.9 mg/dl. That analysis also showed that 24 of 25 patients with acute kidney injury (AKI) had DSS as a final diagnosis. Of the 25 patients with DHF-associated AKI, 16 (64%) died as a result of profound shock, together with other conditions such as liver failure, respiratory failure, and severe bleeding^[37]. Studies in adults have reported an AKI incidence of 14.2% among dengue patients, and those with AKI saw significant morbidity and mortality, longer hospital stays, and poor renal outcomes^[38]. Early diagnosis of dengue infection, known clinical and laboratory characteristics and risk factors together with early detection of AKI using appropriate criteria^[39], and monitoring for warning signs of severe dengue, are essential if AKI and other complications are to be avoided^[40]. Although the two sets of WHO criteria from 1997 and 2009 are still debatable in terms of their ability to appropriately differentiate dengue from OFI and to classify disease severity^[20-22, 41, 42], the problem is compounded by a lack of key data in resource-poor settings, making it difficult to apply the criteria. For instance, we lacked information on clinical bleeding sites, and were only able to detect gastrointestinal bleeding based on stool examinations. In addition, there was a lack of information on blood pressure or narrow pulse pressure to indicate whether a patient was in shock^[43], no data on restlessness suggestive of circulatory failure, and no chest X-ray results to evaluate pleural effusion or abdominal ultrasound to detect ascites, both of which are important for identifying plasma leakage. The 1997 and 2009 WHO dengue guidelines also include a tourniquet test as a diagnostic tool for dengue in the early febrile phase. However, the tourniquet test has been shown to have low sensitivity for dengue diagnosis, such that a negative result does not exclude dengue infection^[44-46]. The tourniquet test had not been performed for the majority of patients in our data set and was thus not included in the analysis. Regarding the two approaches used in this study, CART versus the more conventional approach of logistic regression, some points are worth mentioning here. Firstly, our main focus was on building the decision tree model from CART analysis. CART is non-parametric, and can manipulate numerical data which may be highly skewed, multimodal, ordinal or non-ordinal in structure. CART is not significantly impacted by outliers in the input variables. The output of CART in the form of a decision tree is easy to follow and gives some visual information on the hierarchical importance of the variables from the top to the bottom of the tree, although calculating the importance matrices of the predictors in CART is not straightforward. In this study, therefore, we quantified the importance of each decision tree predictor via the odds ratio as calculated by logistic regression. Secondly, the ways in which the decision boundaries are generated in the two approaches are different. While the logistic regression generates a single boundary, a decision tree essentially partitions the data space into half-spaces using axis-aligned linear decision boundaries, giving a non-linear decision boundary. Either approach may be more applicable depending on the setting.

Finally, the model's accuracy was measured in different ways for each of the two approaches. For the decision tree model, the out-of-sample accuracy was estimated via cross-validation, i.e. the 10-fold cross-validation function in Weka allowed us to conveniently perform the cross-validation and directly re-report the model's accuracy. For the logistic regression, however, the model's accuracy was estimated from the classification table, which showed the number of observed against predicted outcomes, using a default cut-off value of 0.5 for the predicted probability. For all of the above reasons, it would have been difficult to compare the relative merits of the two methods used in our study. There were several limitations with regard to the data-set used for this study. Firstly, the data came from just one hospital, a further indicator of the poor resources in the Southeast Asia region where dengue is endemic. Secondly, due to the lack of IgG antibody results it was not possible to interpret whether cases were primary or secondary dengue infections. This information could potentially be a useful early indicator for the severity of a dengue infection. Thirdly, the algorithm was derived from data collected within 48 h of admission among children aged less than 16 years old. If the model were to be used for older patients or in different regions, some adjustments to it may be necessary. Although the cohort of 198 patients with confirmed dengue was relatively small, with an even smaller subset of just 38 severe dengue cases, the simple model we derived is still likely to be useful because it includes a small number of predictive variables that would probably be available in similar settings. In addition, a previous study by Carter *et al.* showed that the DENV rapid diagnostic test (RDT) had low sensitivity for the diagnosis of dengue infection^[47]. However, the development of diagnostic tests for dengue has advanced rapidly. The NS1 test in particular has become widely available in many resource-limited settings. It is simple to use and has acceptable accuracy. If rapid diagnosis of dengue using NS1 can be achieved, our algorithm would prove very useful. This also highlights the importance of children attending for testing as soon as dengue is suspected, because NS1 detection is optimal during the first seven days of infection. The algorithm will become more relevant and useful as the rapid diagnosis of dengue becomes more common. By using our simple algorithm to help identify and predict severe dengue, we believe that there would be more room to focus on other, more serious bacterial diseases, which are all-too-common in these types of resource-poor settings.

6. Conclusions

Our call tree rule mistreatment straightforward clinical and laboratory indicators contains a moderate classification accuracy for predicting the event of severe break bone fever among pediatric patients with confirmed DENV infection. The model demonstrates the importance of hematocrit and blood platelet levels for observation the severity of dengue fever, as indicated by the WHO criteria and previous studies. Our rule offers straightforward indicators for severity, as well as hematocrit, GCS, excreta macromolecule, creatinine, and blood platelet count, all of that area unit measured on admission. This model is probably helpful for guiding inmate observation and patient management of fever cases. The model will need additional validation against alternative datasets from cohort studies conducted in an exceedingly style of settings, with the goal

of building a universal rule for guiding clinical management of severe dengue fever in resource-limited settings.

Abbreviations

AHC: Angkor Hospital for Children; AKI: Acute kidney injury; ALT: Alanine Aminotransferase; CART: Classification and regression tree; CI: Confidence Interval; DHF: Dengue hemorrhagic fever; DSS: Dengue shock syndrome; GCS: Glasgow Coma Score; ICU: Intensive care unit; JEV: Japanese encephalitis Virus; OFI: Other febrile illnesses; OR: Odds ratio; WBC: White blood cells

7. References

1. Tantawichien T. Dengue fever and dengue haemorrhagic fever in adolescents and adults. Paediatrics and international child health. 2012; 32(Suppl 1):22-7.
2. Tanner L, Schreiber M, Low JG, Ong A, Tolfvenstam T, Lai YL, *et al.* Decision tree algorithms predict the diagnosis and outcome of dengue fever in the early phase of illness. PLoS Negl Trop Dis. 2008; 2(3):e196.
3. Normile D. Tropical medicine. Surprising new dengue virus throws a spanner in disease control efforts. Science (New York, NY). 2013; 342(6157):415.
4. Dwivedi VD, Tripathi IP, Tripathi RC, Bharadwaj S, Mishra SK. Genomics, proteomics and evolution of dengue virus. Briefings in functional genomics. 2017; 16(4):217-27.
5. Guzman MG, Halstead SB, Artsob H, Buchy P, Farrar J, Gubler DJ, Hunsperger E, Kroeger A, Margolis HS, Martinez E, *et al.* Dengue: a continuing global threat. Nat Rev Microbiol. 2010; 8(12 Suppl):S7-16.
6. Manock SR, Jacobsen KH, de Bravo NB, Russell KL, Negrete M, Olson JG, Sanchez JL, Blair PJ, Smalligan RD, Quist BK, *et al.* Etiology of acute undifferentiated febrile illness in the Amazon basin of Ecuador. Am J Trop Med Hyg. 2009; 81(1):146-51.
7. Huy R, Buchy P, Conan A, Ngan C, Ong S, Ali R, Duong V, Yit S, Ung S, Te V, *et al.* National dengue surveillance in Cambodia 1980-2008: epidemiological and virological trends and the impact of vector control. Bull World Health Organ. 2010; 88(9):650-7.
8. Sam SS, Omar SF, Teoh BT, Abd-Jamil J, AbuBakar S. Review of dengue hemorrhagic fever fatal cases seen among adults: a retrospective study. PLoS Negl Trop Dis. 2013; 7(5):e2194.
9. Potts JA, Thomas SJ, Srikiatkachorn A, Supradish PO, Li W, Nisalak A, *et al.* Classification of dengue illness based on readily available laboratory data. Am J Trop Med Hyg. 2010; 83(4):781-8.
10. WHO. World health organization: a global brief on vector-borne diseases. Geneva: World Health Organization, 2014.
11. Duong V, Henn MR, Simmons C, Ngan C, Gavotte L, Viari A, *et al.* Complex dynamic of dengue virus serotypes 2 and 3 in Cambodia following series of climate disasters. Infection, genetics and evolution: journal of molecular epidemiology and evolutionary genetics in infectious diseases. 2013; 15:77-86.
12. National Institute of Statistics, Ministry of Planning Phnom Penh, Cambodia. General Population Census of Cambodia. Cambodia: National Institute of Statistics, Ministry of Planning Phnom Penh, Cambodia; 2008.
13. Vong S, Goyet S, Ly S, Ngan C, Huy R, Duong V, *et al.* Under-recognition and reporting of dengue in Cambodia: a capture-recapture analysis of the National Dengue Surveillance System. Epidemiol Infect. 2012; 140(3):491-9.
14. Breiman L, Friedman J, Stone CJ, Olshen RA. Classification and regression trees, illustrated, reprint, revised edn: Taylor & Francis, 1984, 1984.
15. Chheng K, Carter MJ, Emary K, Chanpheaktra N, Moore CE, Stoesser N, *et al.* A prospective study of the causes of febrile illness requiring hospitalization in children in Cambodia. PLoS One. 2013; 8(4):e60634.
16. Collins AS. Preventing health care-associated infections. In: Hughes RG, editor. Patient safety and quality: an evidence-based handbook for nurses. Rockville (MD): Agency for Healthcare Research and Quality, 2008.
17. Revelas A. Healthcare - associated infections: a public health problem. Niger Med J. 2012; 53(2):59-64.
18. Handbook WHO: Integrated management of childhood illness. World Health Organization, 2005.
19. Blacksell SD. Commercial dengue rapid diagnostic tests for point-of-care application: recent evaluations and future needs? J Biomed Biotechnol. 2012; 2012:151967.
20. Barniol J, Gaczkowski R, Barbato EV, da Cunha RV, Salgado D, Martinez E, *et al.* Usefulness and applicability of the revised dengue case classification by disease: multi-centre study in 18 countries. BMC Infect Dis. 2011; 11:106.
21. Hadinegoro SR. The revised WHO dengue case classification: does the system need to be modified? Paediatrics and international child health. 2012; 32(Suppl 1):33-8.
22. Kalayanarooj S. Dengue classification: current WHO vs. the newly suggested classification for better clinical application? Journal of the Medical Association of Thailand = Chotmaihet thangphaet. 2011; 94(Suppl 3):S74-84.
23. Seni G, Elder J. Ensemble Methods in Data Mining: Improving Accuracy through Combining Predictions. Morgan & Claypool, 2010.
24. Feelders A. Handling missing data in trees: Surrogate splits or statistical imputation? Principles of Data Mining and Knowledge Discovery. Springer Berlin Heidelberg, 1999, 1704.
25. Cevallos Valdiviezo H, Van Aelst S. Tree-based prediction on incomplete data using imputation or surrogate decisions. ELSEVIER, Information Sciences. 2015; 311:163-181.
26. Potts JA, Gibbons RV, Rothman AL, Srikiatkachorn A, Thomas SJ, Supradish PO, *et al.* Prediction of dengue disease severity among pediatric Thai patients using early clinical laboratory indicators. PLoS Negl Trop Dis. 2010; 4(8):e769.
27. Chuansumrit A. Pathophysiology and management of dengue hemorrhagic fever. Blackwell Publishing, 2006, 3-11.
28. Lei HY, Yeh TM, Liu HS, Lin YS, Chen SH, Liu CC. Immunopathogenesis of dengue virus infection. J Biomed Sci. 2001; 8(5):377-88.
29. Hottz E. Platelets in dengue infection. Drug Discovery

- Today. 2011; 8(No. 1-2):e33-e38. Elsevier.
30. Srikiatkachorn A, Gibbons RV, Green S, Libraty DH, Thomas SJ, Endy TP, *et al.* Dengue hemorrhagic fever: the sensitivity and specificity of the world health organization definition for identification of severe cases of dengue in Thailand, 1994-2005. *Clinical infectious diseases: an official publication of the Infectious Diseases Society of America.* 2010; 50(8):1135-43.
 31. Chadwick D, Arch B, Wilder-Smith A, Paton N. Distinguishing dengue fever from other infections on the basis of simple clinical and laboratory features: application of logistic regression analysis. *Journal of clinical virology: the official publication of the Pan American Society for Clinical Virology.* 2006; 35(2):147-53.
 32. Tamibmaniam J, Hussin N, Cheah WK, Ng KS, Muminathan P. Proposal of a clinical decision tree algorithm using factors associated with severe dengue infection. *PLoS One.* 2016; 11(8):e0161696.
 33. The TD. Clinical features of dengue in a large Vietnamese cohort: Intrinsically lower platelet counts and greater risk for bleeding in adults than children. 2012; 6(6):e1679. *PLOS Neglected Tropical Diseases.*
 34. Rao SM. *Internal Medicine Inside.* Herbert Open Access Journals, 2013.
 35. Khan MI, Anwar E, Agha A, Hassanien NS, Ullah E, Syed IA, Raja A. Factors predicting severe dengue in patients with dengue fever. *Mediterranean journal of hematology and infectious diseases.* 2013; 5(1):e2013014.
 36. Vasanwala FF, Puvanendran R, Fook-Chong S, Ng JM, Suhail SM, Lee KH. Could peak proteinuria determine whether patient with dengue fever develop dengue hemorrhagic/dengue shock syndrome?—a prospective cohort study. *BMC Infect Dis.* 2011; 11:212.
 37. Laoprasopwattana K, Pruekprasert P, Dissaneewate P, Geater A, Vachvanichsanong P. Outcome of dengue hemorrhagic fever-caused acute kidney injury in Thai children. *J Pediatr.* 2010; 157(2):3039.
 38. Mallhi TH, Khan AH, Adnan AS, Sarriff A, Khan YH, Jummaat F. Incidence, characteristics and risk factors of acute kidney injury among dengue patients: a retrospective analysis. *PLoS One.* 2015; 10(9):e0138465.
 39. Mallhi TH, Sarriff A, Adnan AS, Khan YH, Hamzah AA, Jummaat F, Khan AH. Dengue-induced acute kidney injury (DAKI): a neglected and fatal complication of dengue viral infection—a systematic review. *J Coll Physicians Surg Pak.* 2015; 25(11):828-34.
 40. Oliveira JF, Burdmann EA. Dengue-associated acute kidney injury. *Clin Kidney J.* 2015; 8(6):681-5.
 41. Gan VC, Lye DC, Thein TL, Dimatatac F, Tan AS, Leo YS. Implications of discordance in world health organization 1997 and 2009 dengue classifications in adult dengue. *PLoS One.* 2013; 8(4):e60946.
 42. Basuki PS, Budiyanto Puspitasari D, Husada D, Darmowandowo W, Ismoedijanto SS, *et al.* Application of revised dengue classification criteria as a severity marker of dengue viral infection in Indonesia. *The Southeast Asian journal of tropical medicine and public health.* 2010; 41(5):1088-94.
 43. WHO. World Health Organization: Dengue guidelines for diagnosis, treatment, prevention and control. In: *Dengue: Guidelines for Diagnosis, Treatment, Prevention and Control, New Edition.* edn. Geneva edn. Geneva: World Health Organization, 2009.
 44. Cao XT, Ngo TN, Wills B, Kneen R, Nguyen TT, Ta TT, *et al.* Evaluation of the World Health Organization standard tourniquet test and a modified tourniquet test in the diagnosis of dengue infection in Viet Nam. *Tropical medicine & international health: TM & IH.* 2002; 7(2):125-32.
 45. Hammond SN, Balmaseda A, Perez L, Tellez Y, Saborio SI, Mercado JC, *et al.* Differences in dengue severity in infants, children, and adults in a 3-year hospital-based study in Nicaragua. *The American journal of tropical medicine and hygiene.* 2005; 73(6):1063-70.
 46. Mayxay M, Phetsouvanh R, Moore CE, Chansamouth V, Vongsouvath M, Sisouphone S, *et al.* Predictive diagnostic value of the tourniquet test for the diagnosis of dengue infection in adults. *Tropical medicine & international health: TM & IH.* 2011; 16(1):127-33.
 47. Carter MJ, Emary KR, Moore CE, Parry CM, Sona S, Putchhat H, *et al.* Rapid diagnostic tests for dengue virus infection in febrile Cambodian children: diagnostic accuracy and incorporation into diagnostic algorithms. *PLoS Negl Trop Dis.* 2015; 9(2):e0003424.