

## Assessment of Diabetes in accordance with Indian Diabetes Risk Score (IDRS) among Adults of Delwara, Rajsamand

Mitin Parmar<sup>1\*</sup>, Milan Chaudhary<sup>2</sup>, Pranav Bhavsar<sup>3</sup>

<sup>1-3</sup> Department of Community Medicine, Ananta Institute of Medical Sciences and Research Centre, Rajsamand, Rajasthan, India

### Abstract

**Introduction:** Diabetes Mellitus (DM) is a metabolic disorder of multiple etiologies characterized by chronic hyperglycemia with disturbance of carbohydrate, protein and fat metabolism resulting from impaired insulin secretion, insulin action, or both. Assessment of the risk of diabetes development acts as a primary prevention measure among the high-risk population.

**Objectives:** To assess the risk of developing Type 2 Diabetes Mellitus using Indian Diabetic Risk Score among adults of Delwara.

**Methods:** A cross sectional study was undertaken among adults of age more than 30 years to assess the risk of development of DM. A pre-validated, structured questionnaire was used to assess the risk of development of DM including the Indian Diabetes Risk Score. The total scores were classified into low, medium and high-risk categories.

**Results:** Mean age of the study population was 46.12±10.9 yrs. Nearly 62% of study population were female, 42.7% had high school education and 44.7% were daily wage employees. Majority (47%) of the study population belonged to high risk and 36.5% belonged to medium risk category for development of DM. Increasing age, BMI, waist circumference, high Blood Pressure, a positive family history, female gender, higher education, current usage of alcohol and tobacco, sedentary to mild physical activities and high waist to height ratio were significantly associated with increased risk of development of Diabetes Mellitus.

**Conclusion:** The risk assessment should be put forward as a major prevention tool in the DM management.

**Keywords:** Risk Assessment, Delwara, Diabetes mellitus.

### 1. Introduction

Diabetes mellitus is a global health menace which is increasing in an alarming rate. The projected global prevalence of diabetes among adults as 9.9% by 2030 from 8.3% in 2011. While communicable diseases are slowly getting controlled in India, there is a significant increase in the burden of non-communicable diseases, including but not restricted to diabetes. The age of man-made and degenerative diseases characterized by a life expectancy close to 50-60 years and unhealthy lifestyles which promote diseases like cardiovascular disease and hypertension and Diabetes Mellitus. The prevalence rate of diabetes mellitus among the adult population ranges from 3% in rural areas to 9% in urban areas in India.

A study by the American Diabetes Association reports that India will see the greatest increase in people diagnosed with diabetes by 2030. The high incidence is attributed to a combination of genetic susceptibility plus adoption of a high-calorie, low-activity lifestyle by India's growing middle class.

Complications of Diabetes Mellitus are mainly acute complications like hypoglycemia, hyperglycemic crisis, diabetic ketoacidosis (DKA), hyperglycemic hyperosmolar state (HHS) and Chronic complications like diabetic retinopathy, diabetic nephropathy, diabetic neuropathy (micro vascular complications) macro vascular complications-cardiovascular diseases such as heart attacks, strokes and insufficiency in blood flow to legs, diabetic foot disease. Other rare complications like impaired growth and development, associated autoimmune conditions,

hypothyroidism, hyperthyroidism, celiac disease, vitiligo, primary adrenal insufficiency (Addison's disease), lipodystrophy (lipoatrophy and lipohypertrophy), non-alcoholic fatty liver disease, infections seen in patients with diabetes, limited joint mobility, edema. These complications are also on the rise and contribute significantly to overall morbidity and mortality. The low levels of education and poor awareness of the disease in the country are enhancing its impact on health of the population.

While comprehensive data are not available, smaller studies have been performed in various states of India to study the prevalence of diabetes. Based on these studies, the highest prevalence reported is from Ernakulum in Kerala (19.5%) and the lowest from Kashmir valley (6.1%). Most other areas have prevalence above 10%. According to National Family Health Survey 4, the prevalence of high blood sugar level among men is 13.1% and women 8.7% in Kerala.

A majority of our population is at high risk of developing Diabetes Mellitus and is not aware of it until complications arise. Our study aims to identify these people and bring about measures to prevent the early onset of Diabetes.

### 2. Materials and Methods

The study was done among adults of age more than 30 years without a history of Type 2 Diabetes Mellitus at Delwara, Rajsamand, Rajasthan, INDIA during a 3 months period. Sample size of 170 was reached using the formula,  $n = z^2pq/d^2$ , Where,  $z$  = relative deviate (at 95% confidence interval) i.e. 1.96,  $p$  = i.e. estimated 36.5% prevalence of high risk of development of DM from a study done in Pune,

**Maharashtra.**

Pretested, structured interview schedule was administered to collect the data after obtaining Institutional Ethics Committee approval. The interview schedule was divided into 2 parts. The first part consisted of socio demographic details of study subjects and risk factors of diabetes. The second part was Indian Diabetic Risk Score which comprised of two modifiable (Waist circumference and Physical activity) and two non-modifiable risk factors (age and family history of T2 DM).

Indian diabetic risk score (IDRS) is the screening tool widely used to assess the risk of diabetes among general population. An IDRS  $\geq 60$  had a sensitivity of 83.8%, specificity of 81.0% in prediction of T2DM, while an IDRS  $< 60$  had a sensitivity of 79.9% and specificity of 83.8%, in prediction of non-T2DM in Indian Settings.

**Operational definitions**

Participants with IDRS  $\geq 60$  were considered at high risk of diabetes. Family history of diabetes If either or both of a subject's parents had diabetes, they were considered to have a positive family history. Physical activity levels were graded based on WHO STEPS definitions of sedentary, mildly, moderately or vigorously physically active. Waist circumference measured to the nearest 0.1 cm at the midpoint between the tip of the iliac crest and the last costal margin in the back and at the umbilicus in the front, using a non-stretchable tape, at the end of normal expiration, with the subject standing erect in a relaxed position. Abdominal/central obesity was considered to be present when the waist circumference was  $\geq 80$  cm in women and  $\geq 90$  cm in men. Weight was measured to the nearest 0.1 kg using a digital weight recorder. Height was measured to the nearest 0.1 cm using a wall fixed stadiometer. Body Mass Index was evaluated as weight divided by height squared ( $\text{kg}/\text{m}^2$ ). Waist to Height Ratio was calculated as waist circumference divided by height. The data collected was entered in Microsoft Excel and analyzed using Statistical Package for Social Sciences version 16 software. Descriptive analysis was done by calculating frequencies, proportions, mean, median, standard deviations. Chi square tests used to calculate associations.

**3. Results**

Mean age of the study population was  $46.12 \pm 10.9$  yrs. Among 170 study population 87 (51.2%) belonged to the age group of 35-49 years. Nearly 67% of study population were female, 42.7% had high school education. About 39 (23%) and 41 (24%) were currently using alcohol and tobacco respectively. Nearly 98% were following non-vegetarian diet. 32 (18.8%) of study subjects had associated hypertension and on treatment. About 58% of study subjects have family history of Diabetes. Mean waist to height ratio was  $0.61 \pm 0.11$ . Majority 149 (87.6%) had a high waist to height ratio. 44 (25.9%) of study subjects had raised blood pressure during the interview.


Majority (90%) had a family size  $< 5$  and nearly 62% belonged to nuclear family. About 70% of females were homemaker and 44.1% of study population was daily wagers. Nearly 56% of study population didn't have any comorbidity. 32 (18.8%) had associated hypertension and 23 (13.5%) had osteoarthritis, 20 (11.7%) had others like asthma, stroke, Acid Peptic Disease, migraine. (Figure 1) About 80 (47%) of the study population belonged to high

risk and 36.5% belonged to medium risk category for development of Diabetes Mellitus according to Indian Diabetic Risk Score (IDRS). (Table 1)

Increasing age, BMI, waist circumference, high Blood Pressure, a positive family history and female gender had a strong association with high risk of development of diabetes. We found that higher the education higher the risk for Diabetes. Current usage of alcohol and tobacco were also associated with higher risk of development of Diabetes Mellitus. Sedentary to mild physical activities and high waist to height ratio were also associated with increased risk of development of Diabetes Mellitus.

**Table 1: Indian Diabetes Risk Score**

IDRS	Frequency (%)
Low Risk	28 (16.5)
Medium Risk	62 (36.5)
High Risk	80 (47)



**Fig 1: Associated medical conditions**

**4. Discussion**

In this study we had found that majority of our study population belonged to age group 36-49 years, nearly 62% of study population was female, 42.7% had high school education and 44.7% were daily wage employees. This was similar to a study done in Maharashtra. Using the IDRS we found that majority of 36-49 years age group belonged to moderate risk and in age group of  $> 50$ , majority was under high risk of development of Type 2 Diabetes Mellitus. Many studies conducted in India showed that increasing age increases the risk of Diabetes. In another study which used Diabetes Risk Score, the age group of 55-64 yrs was high risk of developing Diabetes.

In our study nearly 58% of overweight study subjects and 71% of obese people had a high risk of development of Diabetes. Studies have shown that obesity plays a major factor in the development of diabetes. Obesity and diabetes are interrelated and the term 'Diabesity' has been coined to describe the relationship between obesity and diabetes.

Another significant association was waist circumference with risk of developing Diabetes. Our study showed waist circumference and risk of developing Diabetes is directly proportionate as increased waist circumference clearly denote deposit of fat around internal organs leading to insulin resistance. Other studies showed similar results.

Among our population with high BP, about 77.3% were in the high-risk group of developing Diabetes. Diabetes damages arteries and makes them targets for hardening and atherosclerosis leading to hypertension. A population-based study done in Chennai, India among the NGT (normal glu-

cose tolerance) subjects showed about 38.2% with Hypertension came under the high-risk group of IDRS. Apart from obesity the other risk factors for type 2 DM are age, life style factors like sedentary life style, smoking and alcohol consumption and the most important risk factor was family history of type 2 dm which has a critical role in the development of the disease which is similar to a study done in Chennai.

We did not find any significant association b/w socioeconomic status and risk of developing Diabetes. A cross-sectional study conducted at Pune, Maharashtra showed that the association between socioeconomic class and risk status was highly significant statistically ( $P = 0.001$ ). We did not find any association between type of diet, occupation, family income and risk of developing Diabetes.

Every adult more than 18 years should be assessed for risk of development of Diabetes Mellitus using IDRS or any other simple tool. It will not just aid in early detection and prevention of complication of Type 2 DM but also helps to reduce the morbidity and mortality associated with it. This tool can be used as a primary prevention measure to motivate people against metabolic diseases and cardiovascular diseases. As majority of study population belonged to high and moderate risk for development of Diabetes, urgent follow up and health education are needed in the area with a Random Blood Sugar testing for all followed up by definitive diagnostic tests. Development of comprehensive health approaches including dietary and lifestyle modifications and awareness regarding the risk of Diabetes Mellitus and its complications should be done.

The assessment of risk would have been complete if qualitative component also added. That would have shed a light regarding the resistance towards adapting the healthy lifestyle.

## 5. Conclusion

Majority (47%) of the study population belonged to high risk and 36.5% belonged to moderate risk category for development of Diabetes Mellitus. Increasing age, BMI, waist circumference, high Blood Pressure, a positive family history and female gender had a strong association with high risk of development of diabetes. We found that higher the education higher the risk for Diabetes. Current usage of alcohol and tobacco were also associated with higher risk of development of Diabetes Mellitus. Sedentary to mild physical activities and high waist to height ratio were also associated with increased risk of development of Diabetes Mellitus.

## 6. References

- Whiting DR, Guariguata L, Weil C, Shaw J. IDF diabetes atlas: global estimates of the prevalence of diabetes for 2011 and 2030. *Diabetes Research and Clinical Practice*. 2011; 94(3):311-21.
- Ramachandran A, Snehalatha C, Shyamala P, Vijay V, Viswanathan M. High prevalence of NIDDM and IGT in an elderly south Indian population with low rates of obesity. *Diabetes Care*. 1994; 17(10):1190-2.
- Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: Estimates for the year 2000 and projections for 2030. *Diabetes Care*. 2004; 27(5):1047-53.
- Benja M, *et al*. Use of glucometer and fasting blood glucose as screening tools for diabetes mellitus type 2 and glycated haemoglobin as clinical reference in rural community primary care settings of a middle-income country. *BMC Public Health*. 2012; 12(1):349.
- Sankar VU, Kutty RV, Santhosh KN. Life Course Socioeconomic Transition and its Association with Early Onset Type 2 Diabetes: Protocol for a Sequential Exploratory Mixed Method Study. *J Clin Diagn Res*. 2016; 10(6):LO01-5.
- "Diabetes Programme". World Health Organization. Archived from the original on 26 April 2014, 2014.
- Raman KV, Aleyamma J, Soman CR. High Prevalence of Type 2 Diabetes in an Urban Settlement in Kerala, India. *Ethn Health*. Published online. 1999; 4(4):231-9
- Vasudevan U, Selvamani Y. Rise of Health Risks among Adults in India: A Quick Look at NFHS-4 Fact Sheets. *Social Science Spectrum*. 2016; 1(4):324-7.
- Patil RS, Gothankar JS. Assessment of risk of type 2 diabetes using the Indian Diabetes Risk Score in an urban slum of Pune, Maharashtra, India: a cross-sectional study. *WHO South-East Asia J Public Health*. 2016; 5(1):53-61.
- Mohan V, Sandeep S, Deepa M, Gokulakrishnan K, Datta M, Deepa R. A diabetes risk score helps identify metabolic syndrome and cardiovascular risk in Indians – the Chennai Urban Rural Epidemiology Study (CURES-38). *Diabetes, Obesity and Metabolism*. 2007; 9(3):337-43.
- Sharma KM, *et al*. Indian Diabetes Risk Score helps to distinguish type 2 from non-type 2 diabetes mellitus (GDRC-3). *J Diabetes Sci Technol*. 2011; 5(2):419-25.
- Shah B. Development of Sentinel Health Monitoring Centres for Surveillance of Risk Factors of Non communicable Diseases in India (April 2003 to March 2005): Collated Results of 6 Centres. *Indian Council of Medical Research: New Delhi*, 2005.
- Rao CR, Kamath VG, Shetty A, Kamat A. A study on the prevalence of type 2 diabetes in coastal Karnataka. *Int J Diabetes Dev Ctries*. 2010; 30(2):80-5. doi:10.4103/0973-3930.62597.
- Thirunavukkarasu S, Srinivasan K, Sankara SP, Kavumpurathu RT. Achutha Menon Centre Diabetes Risk Score: A Type 2 Diabetes Screening Tool for Primary Health Care Providers in Rural India. *Asia Pac J Public Health*. 2015; 27(2):147-54.
- Mohan V, Sandeep S, Deepa M, Gokulakrishnan K, Datta M, Deepa R. A diabetes risk score helps identify metabolic syndrome and cardiovascular risk in Indians – the Chennai Urban Rural Epidemiology Study (CURES-38). *Diabetes, Obesity and Metabolism*. 2007; 9(3):337-43.
- Anjana RM, Lakshminarayanan S, Deepa M, Farooq S, Pradeepa R, *et al*. Parental history of type 2 diabetes mellitus, metabolic syndrome and cardio metabolic factors in Asian Indian adolescents. *Metabolism Clinical and Experimental*. 2009; 58(3):344-50.
- Padaki S, Vijayakrishna K, Dambal A, *et al*. Anthropometry and physical fitness in individuals with family history of type 2 Diabetes mellitus: A comparative study. *Indian Journal of Endocrinology and Metabolism*. 2011; 15(4):327-30.