



Co-relation between cervicothoracic angle and neck pain in adults

Amit Dwivedi¹, Vikram Dagar^{2*}, Shivani Tiwari Dwivedi³, Sonam⁴, Fenil Shah⁵

¹ Associate Professor, Department of Orthopedics, Santosh Medical College & Hospital, Ghaziabad, Uttar Pradesh, India

² Postgraduate Student, Department of Orthopedics, Santosh Medical College & Hospital, Ghaziabad, Uttar Pradesh, India

³ Consultant, Lal Path Labs, Ghaziabad, Uttar Pradesh, India.

⁴ Junior Resident, Sharda hospital, greater noida, Uttar Pradesh, India.

⁵ Postgraduate Student, Department of Orthopedics, Santosh Medical College & Hospital, Ghaziabad, Uttar Pradesh, India

* Corresponding Author: Vikram Dagar

Abstract

Introduction: Neck pain is the fourth most common cause of disability after lower back pain, depression, and joint pain. Cervical sagittal balance is as crucial as pelvic sagittal alignment and is related to the concept of T1 alignment.

Methods and Materials: 235 Patients diagnosed as neck pain and treated at our institute between August 2017 to July 2019 with age between 20-80 years with neck pain complaints and on medication were included in this study. Pain and functional improvements were assessed using visual analogue scale (VAS) and neck disability index (NDI). Standing lateral view and standing swimmers lateral view of cervical spine radiographs were taken and studied for evaluating cervicothoracic parameters T1 slope and SVA (Sagittal Vertical Axis) C2-7, following neck pain and compared with normal ranges. Variations of these criteria have been reported along with the scores of the questionnaire. Statistical analysis was carried out using the edition 21.0 of the Statistical Package for Social Sciences (SPSS).

Results: Our analysis found that the average T1 slope was $27.82 + 14.33$, the average male T1 slope was $26.74 + 14.21$ and the average female T1 slope was $28.56 + 14.42$. According to Sang *et al*⁷ average T1 slope is $25.7.5 + 6.4$ which was taken as a reference for comparison with the asymptomatic population, our study had an increased value but was not significant.

Conclusions: The pain in the neck increases with age. It is more prevalent in females. Study shows an increase in neck pain with increasing age due to degenerative changes in the T1 slope, SVA C2-C7. There is no significant correlation with cervical and neck pain or disability but a good relationship between the two. There was no substantial difference in cervical curve between symptomatic and asymptomatic patients.

Keywords: neck pain, cervical pain, cervicothoracic junction, T1 slope

Introduction

Neck pain is the fourth most common cause of disability after lower back pain, depression, and joint pain^[1]. Simple conditions that are prone to accumulation such as mechanical stress, lack of muscle strength, office computer jobs, non-ergonomic working environments, and long working hours result in neck pain being more frequently seen in middle age^[2].

Cervical vertebral X-ray is the most common diagnostic tool used in this circumstance^[3, 4]. Cervical sagittal balance is as crucial as pelvic sagittal alignment and is related to the concept of T1 alignment^[5].

Due to a lack of clear consensus on the relationship between cervico-thoracic parameters and neck pain in the literature, we aimed to determine the various cervical thoracic junction parameters of our patients with radiographs due to better affordability.

Aim

The aim of this present study is aimed to examine whether the sagittal profile of the cervicothoracic spine shows any association with the presence and severity of neck pain in the adult population.

Material Methods

Study Design: Prospective observational study.

Place of Study: Patients attending the Out-patient Department of Orthopaedics, Santosh Medical College and Hospital, Ghaziabad, Uttar Pradesh, India.

Duration of Study: Study was conducted between August 2017 to July 2019.

Inclusion Criteria • Age between 20-80 years. • Patients with neck pain complaints. • Patient already on medication.

Exclusion Criteria • Congenital cervical spinal deformity. • Any motor and sensory deficits. • Significant previous or recent trauma to the spine. • Previous cervical spinal surgery. • Spinal tumors and metastasis. • Significant history of malignancy or family history of malignancy. • Infections of spine. • Presence of spinal deformity on forward bending test. • Metabolic bone diseases. • Contraindications to radiographs (e.g., existing or suspected pregnancy). • Presence of red flag symptoms like constant pain, night pain, fever, loss of weight, and loss of appetite. • Malingers. • Patient refusing consent.

Sample Size

The study of Askin A, *et al.* [6] observed correlation between cervical angle and NDI was -0.095. Taking this value as reference, the minimum required sample size with 90% power of study and 5% level of significance is 230 patients. To reduce margin of error, total sample size taken is 235.

Approval obtained from the Ethical Committee of Santosh University and consent taken from each participant.

Patient’s information, history, pre-tested pain scale, and pre-tested disability index were obtained. The patients were explained in detail about the condition and the study protocol and informed consent were taken for utilizing the radiograph and examination findings for the research work. Standing lateral view and standing swimmers lateral view of cervical spine radiographs were taken and studied for evaluating various cervicothoracic parameters following neck pain and compared with normal ranges of the parameter studied.

Radiographic Examination following clinical assessment by the attending specialist, all patients underwent standing lateral view and standing lateral swimmers view of cervical spine. Measurement of Radiographic Parameters The radiographic film cassette was placed 72 inches from the tube, and radiographs were taken without magnification. Digital X-ray photographs were collected from the PACS method, which were used to collect orientation measurements.



Fig 1

Standing lateral view and standing swimmers lateral view of cervical spine radiographs were taken and studied for evaluating cervicothoracic parameters T1 slope and SVA C2-7, following neck pain and compared with normal ranges of the parameter studied. Variations of these criteria have been reported along with the scores of the questionnaire. Statistical analysis was carried out using the edition 21.0 of the Statistical Package for Social Sciences (SPSS).

Observations and Results

In this research, 13.6 percent of patients in the age group between 20 and 40 years of age, 58.4 percent of patients in the age group between 41 and 60 years of age, and 28.1 percent in the age group between 61 and 80 years of age. Most patients had dominant arm pain patterns. The average NDI was 56.54 + 12.84, the average male NDI was 57.2 +

12.43 and the average female NDI was 56.1 + 13.14. The average VAS was 5.9 +2.17, the average male VAS was 5.7 + 2.31 and the average female VAS was 5.92 + 2.06. Total T1 slope was 27.82 + 14.33, the mean male T1 slope was 26.74 + 14.21 and the mean female T1 slope was 28.56 + 14.42.

Average SVA C2-C7 was 18.55 + 12.75, Average male 17.34 + 11.99 and Average female 19.36 + 13.25. The T1 slope in patients with neck pain is 27.82 and the T1 slope in asymptomatic patients is 25.7 (p=0.2098). The SVA C2-C7 in patients with neck pain is 18.55 and the SVA C2-C7 in asymptomatic patients is 17 (p=0.60255).




Fig 2

Table 1

Variable	Neck pain	Asymptomatic	P-value
T1 slope	27.82	25.7	0.2098
SVA C2-C7	18.55	17	0.60255




Fig 3

Discussion

Our analysis found that the average T1 slope was 27.82 + 14.33, the average male T1 slope was 26.74 + 14.21 and the average female T1 slope was 28.56 + 14.42. According to Sang *et al*⁷ average T1 slope is 25.7.5 + 6.4 which was taken as a reference for comparison with the asymptomatic population, our study had an increased value but was not significant.

In our study average SVA C2-C7 was 18.55 + 12.75, average male 17.34 + 11.99 and average female 19.36 + 13.25. Guo *et al*⁸ also found out that asymptomatic population average SVA C2-C7 (mm) was 16.9 + 10.6. Our study had no significant difference but slightly higher. Based on limited data that could be obtained under the eligibility criteria of this study, there was no significant

difference between symptomatic and asymptomatic individuals in the angle of lordosis of the cervical spine. However, this observation is identical to several other associated studies.

Conclusion

In this research, we found that the pain in the neck increases with age. It is more prevalent in females. Study shows an increase in age due to degenerative changes in the T1 slope, SVA C2-C7. There is no significant correlation between cervical and neck pain or disability but a good relationship between the two. There is no substantial difference in cervical curve between symptomatic and asymptomatic patients.

There is therefore an immediate need for regular physical inspection, calcium and vitamin D supplements and regular exercise.

As our sample size was small, it stopped us from obtaining more conclusive results.

Abbreviations

SVA – Saggital Vertical Axis

VAS – Visual Analogue Scale

NDI – Neck Disability Index

References

1. Murray JL, *et al.* US Burden of Disease Collaborators. The state of US health, 1990- 2010: burden of diseases, injuries, and risk factors. *JAMA*, 2013; 310:591-608.
2. Yang H, Haldeman S, Nakata A, Choi B, Delp L, Baker D, *et al.* Work-related risk factors for neck pain in the US working population. *Spine (Phila Pa 1976)*, 2015; 40:184-192.
3. Gore D, Sepic S, Gardner G. Roentgenographic findings of the cervical spine in asymptomatic people. *Spine*, 1986; 11:521-524.
4. Grob D, Frauenfelder H, Mannion AF. The association between cervical spine curvature and neck pain. *Eur Spine J*, 2007; 16:669-678.
5. Brattberg G, Thorslund M, Wikman A. The prevalence of pain in a general population. The results of a postal survey in a county of Sweden. *Pain*. 1989; 37:215-22.
6. Aşkin A, *et al* "The evaluation of cervical spinal angle in patients with acute and chronic neck pain.": *Turk J Med Sci*. 2017; 47(3):806-811.
7. Sang, *et al.* Factors determining cervical spine sagittal balance in asymptomatic adults: correlation with spinopelvic balance and thoracic inlet alignment": *The Spine Journal*, 2013.
8. Guo, *et al.* "Cervical lordosis in asymptomatic individuals: a meta- analysis": *Journal of Orthopaedic Surgery and Research*, 2018, 13:147.