



A clinical and demographic profile of atrial fibrillation-A hospital based descriptive study

Sourav Chattopadhyay¹, Aniket Sinha^{2*}, Suman Kumar Singh², Abhinav Choudhary², Sumit Singh Kaushal²

¹ Associate Professor, Department of General Medicine, M.G.M. Medical College and L.S.K. Hospital, Kishanganj, Bihar, India

² Resident, Department of General Medicine, M.G.M. Medical College and L.S.K. Hospital, Kishanganj, Bihar, India

Abstract

Background: Atrial Fibrillation (AF) is an abnormal heart rhythm characterized by the rapid and irregular beating of the atrial chambers of the heart. High blood pressure and valvular heart disease are the most common alterable risk factors for AF. The aim of this study was to evaluate the clinical profile and echocardiographic features in patients of AF.

Methods: A total of 110 patients aged above 18 years and diagnosed with AF were enrolled in the study in MGM Medical College & LSK Hospital, Kishanganj, Bihar. These patients underwent a detailed general physical and systemic examination. Relevant biochemical investigations, chest X-ray and echocardiography were done in all patients.

Results: The mean age of the study group was found to be 62.8 years with a male preponderance. Hypertension was the most common cause overall contributing to 28.79 % followed by rheumatic heart disease (21.2 %). A total of 36 patients exhibited LA size greater than 5cm and LA clot was observed in 4% of patients on transthoracic echocardiography.

Conclusion: Hypertension was seen to be a major risk factor for AF and hence it needs to be detected and managed early. Lifestyle modifications with diet and exercise can help in reducing the prevalence of hypertension and thus the risk of developing AF.

Keywords: demographic profile, hospital based descriptive

Introduction

Atrial fibrillation is the most common type of cardiac arrhythmia. It is due to abnormal electrical activity within the atria of the heart, causing them to fibrillate. It is characterized as a tachyarrhythmia, which means that the heart rate is often fast. This arrhythmia may be paroxysmal (less than seven days) or persistent (more than seven days). Due to its rhythm irregularity, blood flow through the heart becomes turbulent and has a high chance of forming a thrombus (blood clot), which can ultimately dislodge and cause a stroke. Atrial fibrillation is the leading cardiac cause of stroke. Risk factors for atrial fibrillation include advanced age, high blood pressure, underlying heart and lung disease, congenital heart disease, and increased alcohol consumption. Symptoms vary from asymptomatic to symptoms such as chest pain, palpitations, fast heart rate, shortness of breath, nausea, dizziness, diaphoresis (severe sweating), and generalized fatigue. Although atrial fibrillation may be a permanent disease, various treatments and risk modifying strategies have been developed to help reduce the risk of stroke in patients that remain in atrial fibrillation. Treatments include anticoagulation, rate control medication, rhythm control medication, cardioversion, ablation, and other interventional cardiac procedures.

Materials and Methods

The present study was an observational time bound study conducted from 2020-2021. All the patients who presented with features suggestive of AF and were 18 years or older were enrolled in the study in MGM Medical College & LSK Hospital, Kishanganj, Bihar. A diagnosis of AF was confirmed by electrocardiographic (ECG) changes. A written informed consent was taken from all the participants. A detailed history with chief complaints of dyspnea, chest pain, palpitations and fatigue were obtained. An enquiry was made into the associated co-morbid conditions and personal habits of patients. All patients underwent elaborate general and systemic examination. The cardiovascular system examination included detailed assessments such as inspection, palpation, and auscultation of cardiovascular system, presence of murmurs, looking for signs of cardiac failure and pulmonary artery hypertension (PAH), and the presence of cardiomegaly. All patients underwent routine investigations Complete blood count, renal function tests, Liver function tests, Chest X-ray, Thyroid profile and ECG. AF was diagnosed based on the ECG findings of absence of P waves, irregular R-R intervals with a rapid

ventricular rate. A 2D-Echocardiogram (2D-ECHO) was done in all ECG proven AF patients, to determine LA (Left atrial) and LV (Left ventricular) dimensions; assess the LV wall thickness and function as well as to recognize the underlying pathological conditions so as to know the etiology of AF (valvular, myocardial or pericardial etiology). All patients were evaluated using M Mode and Two-dimensional transthoracic echocardiography and colour flow Doppler examination. Each 2D ECHO consisted of parasternal long and short axes, and apical two and four chamber views. The conventions of American Society of Echocardiography were followed in obtaining LA dimensions, LV end systolic and end diastolic dimensions, as well as right ventricular end diastolic dimensions. The anterior-posterior LA dimension was calculated in the short axis view from M-mode echocardiography and LA area or volume was measured in apical-chamber or fourchamber views. Left ventricular volumes were measured and ejection fraction (EF) was calculated.

Results

Patient Demographics

A total of 110 patients above 18 years diagnosed with a new onset AF were included in the study. The mean age of the study group was 62.8 years with a male to female ratio of 1.5:1. Majority of the patients (40 %) were in the age group of 46-65 years. Table 1 illustrates the age distribution of patients with AF.

Presenting Clinical Features

Majority of the patients (n=66) presented with dyspnea followed by pedal oedema (n=25). The other presenting features were fatigue, stroke, palpitation, chest pain, syncope and giddiness. Out of 110 patients, 35 gave a history of smoking and 9 patients had history of alcohol consumption. An elevated jugular venous pulse (JVP) was found in 20 % of cases, an irregularly irregular pulse was observed in 90 % of cases. A palpable P2 was observed in 39 patients. A varying S1 was found in 83 % and a loud P2 was heard in 19 % on auscultation. Pansystolic murmur was heard over both mitral and tricuspid area in 11 and 42 patients, respectively and a mid-diastolic murmur was heard in 16 patients. Presystolic accentuation was noticed in 9 % of patients. Around 20 % of patients had crepitations on auscultation suggestive of pulmonary edema. Neurological deficits were detected in 6 %, and heart failure in 15 % of patients. The type of AF commonly encountered in the present study was the persistent type (n=67), followed by paroxysmal (n=26) and permanent (n=17).

Etiology of AF

The commonest cause of AF seen in our study was from non-valvular group contributing to 75.9% of cases. Hypertension was observed to be the most common non-valvular cause for AF. The other non-valvular causes of AF is depicted in Table 2. Amongst the valvular etiology of AF, rheumatic heart disease (RHD) was the commonest cause accounting for 25 out of 110 patients. The valvular involvement in RHD is represented in Table 3.

Investigation Findings

The findings on chest X-ray included cardiomegaly (32 %), COPD (21 %), pulmonary oedema (19 %) and 5 % had acute respiratory distress syndrome (ARDS) due to acute febrile illness and the remaining 23% revealed no findings on chest Xray. All 110 patients were found to have absence of p wave with a varying R-R interval. However, 95 patients had additional ECG findings such as non-specific ST-T changes (33 %), left ventricular hypertrophy (LVH) (21 %), right ventricular hypertrophy (RVH) (18 %), right bundle branch block (RBBB) (12%), left bundle branch block (LBBB) (5 %), AF with aberrancy and block (4 %) and digitalis effects (3 %). The commonest ECHO findings noted in this study were pulmonary artery hypertension (59 %), tricuspid regurgitation (58 %), RHD (21 %), Dilated atria (40 %), Systolic dysfunction (9 %), LVH (29 %) and regional wall motion abnormalities suggestive of IHD (9 %). Left atrial clot was noticed in 4% of patients.

Table 1: Age Distribution of Patients with AF

Age in Years	No. of Patients
18-45	16
46-65	43
66-75	30
>75	21

Table 2: Non Valvular Causes of AF

Etiology	Number(n=83)
Hypertension	32
COPD	19
Diabetes Mellitus	11
IHD	7
Dilated Cardiomyopathy	5
Thyrotoxicosis	4
Lone AF	1

Table 3: Valvular involvement in RHD

Valvular involvement	Number(n=27)
MS with TR	10
MS with MR with TR	8
Isolated MS	6
MR	3

Table 4: Comparison of Risk Factors and Associations of AF with Earlier Studies

	The Framingham Study	Cardiovascular Health Study	ALFA Study	Present
Mean Age	72	-	68.7	62.8
Smoking	34.8%	-	-	32.03%
DM	16.3%	-	10.7%	14.1%
HTN	44%	-	39.4%	28.79%
CHF	20.6%	19%	29.8%	16%
RHD	16%	4.3%	15.2%	21.2%
IHD	25%	22%	16.6%	6.7%
Stroke	-	14.8%	8.4%	5%
Thyrotoxicosis	-	2.4%	3.1%	3.7%
COPD	36%	7.7%	11.2%	17.7%

Discussion

Atrial fibrillation is the most common arrhythmia encountered in clinical practice characterized by uncoordinated atrial activity resulting in an irregularly irregular ventricular rhythm with a rate of 300-600 beats per minute.² It has an estimated prevalence of 3 % in adults and its prevalence increased progressively with age. Some of the well-known risk factors of AF include hypertension, valvular heart disease, male gender, obesity and alcohol consumption.³ The mean age of presentation in the present study was found to be 62.8 years with majority of the patients above 45 years of age. This was in concordance with the study by Mohan G *et al* wherein the mean age was 67.02 ± 12.50 years with more than 90 % of patients above 50 years of age.² The prevalence of atrial fibrillation increases substantially with age. This was an observed trend in several other studies, the most notable of them being the Rotterdam study and ATRIA study^[4,5]. However, these studies have been limited by relatively small numbers of patients with AF, restriction of age ranges, ethnic homogeneity.⁶⁻⁹ There was a slight male preponderance in our study similar to other earlier studies. Men are more susceptible to the development of AF. As prevalence of diabetes, hypertension, COPD, and IHD is more common in men and differences in atrial effective refractory period (ERP) in response to rapid atrial pacing have been reported in men and women. The etiological factors of AF were categorized into valvular and non-valvular groups. Non-valvular causes formed the majority in the present study (76.7 %) out of which hypertension formed a significant majority (37.9%). This was similar to the study by Mohan G *et al*, however, this finding did not correlate with the study by Singh *et al* who reported valvular etiology (RHD) as the cause in 60% of patients.^{2, 10} In the present study, valvular cause in the form of RHD was observed in 23.3 % of patients. Risk factors for developing AF in the Framingham Heart study were hypertension (44 %), diabetes (16 %), prior myocardial infarction (26 %), COPD (36 %) and valvular heart disease (17 %).¹¹ Similar findings have been reported in the Cardiovascular Health Study as well.^[12] Increase in LV thickness and stiffness due to hypertension impairs LV diastolic function which leads to LA stretch and pressure and LA remodeling and dysfunction. Structural, contractile, architectural and electrophysiological changes in atria collectively called atrial cardiomyopathy occurs due to atrial remodeling ultimately leading to AF.¹³ The other risk factors under non-valvular group include COPD (17.7 %), diabetes mellitus (14.1 %), IHD (6.7 %), dilated cardiomyopathy (6.3 %), and thyrotoxicosis (3.7 %). Neurological deficits in the form of stroke was observed in 5% of our patients and 16 % of our patients showed features of heart failure. Table 4 illustrates the comparison of the present study with earlier studies conducted.^{12, 14} Majority of our patients (62.1 %) presented with dyspnea followed by pedal edema (23.3 %). Similarly, Mohan G *et al* reported dyspnea and pedal edema as the predominant symptoms in 56% and 39% of patients, respectively.² However, Shaik Sulaiman Meeran *et al* reported palpitation and dyspnea as the predominant symptoms in 52 % and 14 %, respectively.³ Similarly, ALFA study, reported palpitations in 54% of the patients, dyspnea in 44%, angina in 10% and syncope and dizzy spells in 10 %.¹⁴ Palpitations on exertion was observed in 11 % of patients in the present study. Echocardiography (ECHO) is a vital tool in the diagnostic armamentarium of AF as it gives valuable information regarding the cardiac function, complications and prognosis in AF.¹⁵ Left atrial dilatation indicates chronicity and left ventricular dysfunction, thus resulting in AF. Normally LA size is less than 40mm, however, size more than 45-50mm increases the risk of development of AF by four-fold and it is observed that sustained AF in turn can further increase the LA size. Thus, LA size is an important predictor of AF as well as its complications such as stroke as 20-30 % of strokes occur as a result of AF.¹⁶ In the present study 32 % of patients were found to have a LA size of more than 50 mm. In addition, a study by Mahmood ul Hassan *et al* reported a significant correlation between LA size of greater than 45mm and LA clot in AF.^[17] LA clot was observed in only 4% of patients in the present study

unlike the study by Mohan G *et al* with 14.5 % of patients exhibiting LA clot. It is noticed that trans esophageal ECHO has a greater sensitivity in picking up LA clots in comparison with a transthoracic ECHO. [18] This lower percentage could be attributed to the fact that all the patients in the present study were subjected to transthoracic ECHO, hence, the presence of LA clot could have been easily missed on transthoracic ECHO.

Conclusion

Hypertension is one of the leading causes of AF, accounting for most of the cases of AF. In the present study, RHD is the second most common cause, its prevalence reducing owing to the rampant usage of antibiotics. As the prevalence of hypertension is steadily on the rise, it contributes significantly to the cardiovascular risk and death. However, adopting healthy lifestyle habits, early screening and control of hypertension can in turn reduce the risk of developing AF.

References

1. Michaud GF, Stevenson WG. Atrial Fibrillation. In: Jameson J, Fauci AS, Kasper DL, Hauser SL, Longo DL, Loscalzo J. eds. Harrison's Principles of Internal Medicine, 20e New York, NY: McGraw-Hill.
2. Mohan G, Kaur S, Kaur R, Aggarwal A. The study of clinical and echocardiographic assessment of patients with atrial fibrillation. *Int J Adv Med*,2019;6:1041-5.
3. Shaik Sulaiman Meeran, Agila Saravanan M, Geetha S, Sangeetha SV, Krishnakumar S. A study on clinical profile of patients with atrial fibrillation. *IAIM*,2018;5(2):105110.
4. Allesie MA, Boyden PA, Camm AJ *et al*. Pathophysiology and prevention of atrial fibrillation. *Circulation*,2001;103:769-777.
5. Ott A, Breteler MM, de Bruyne MC, van Harskamp F, Grobbee DE, Hofman A. Atrial fibrillation and dementia in a population-based study: the Rotterdam Study. *Stroke*,1997;28(2):316-21.
6. Go AS, Hylek EM, Phillips KA *et al*. Prevalence of Diagnosed Atrial Fibrillation in Adults National Implications for Rhythm Management and Stroke Prevention: the AnTicoagulation and Risk Factors In Atrial Fibrillation (ATRIA) Study. *JAMA*,2001;285(18):2370-2375.
7. Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation as an independent risk factor for stroke: the Framingham Study. *Stroke*,1991;22:983-988.
8. Lake FR, Cullen KJ, de Klerk NH, McCall MG, Rosman DL. Atrial fibrillation and mortality in an elderly population. *Aust N Z J Med*,1989;19:321-326.
9. Phillips SJ, Whisnant JP, WM OF, *et al*. Prevalence of cardiovascular disease and diabetes mellitus in residents of Rochester, Minnesota. *Mayo Clin Proc*,1990;65:344-359.
10. Singh R, Kashyap R, Bhardwaj R, Marwaha R, Thakur M, Singh TP *et al*. The clinical and etiological profile of atrial fibrillation after echocardiography in a tertiary care centre from North India - a cross sectional observational study. *Int J Res Med Sci*,2017;5:847-50.
11. Al-Khatib SM, Wilkinson WE, Sanders LL, McCarthy EA, Pritchett EL. Observations on the transition from intermittent to permanent atrial fibrillation. *American heart journal*,2000;140(1):142-5.
12. Benjamin EJ, Levy D, Vaziri SM, D'agostino RB, Belanger AJ, Wolf PA. Independent risk factors for atrial fibrillation in a population-based cohort: the Framingham Heart Study. *Jama*,1994;271(11):840-4.
13. Prystowsky EN, Padanilam BJ, Fogel RI. Treatment of atrial fibrillation. *JAMA*,2015;314(3):278-288
14. Levy S, Maarek M, Coumel P, Guize L, Lekieffre J *et al*. for the College of French Cardiologists. Characterization of different subsets of atrial fibrillation in general practice in France: the ALFA study. *Circulation*,1999;99:3028-5.
15. Troughton RW, Asher CR, Klein AL. The role of echocardiography in atrial fibrillation and cardioversion. *Heart*,2003;89(12):1447-54.
16. HK Chopra, GS Wander, Praveen Chandra, Viveka Kumar. Atrial Fibrillation Update – A textbook of Cardiology, First edition, 2017, 262263.
17. ul Hassan M, Hussain C, Gul AM, ullah Jan H, Hafizullah M. Frequency of left atrial and appendage clot in patients with severe mitral stenosis. *J Ayub Med College Abbottabad*,2010;22(2):40-2.
18. Kapral MK, Silver FL. with Canadian Task Force on Preventive Health Care. Preventive health care, 1999 update: 2. Echocardiography for the detection of a cardiac source of embolus in patients with stroke. *CMAJ*,1999;161:989-996